

Southern United States Soybean Disease Loss Estimates for 2023

Allen, T.W.¹, Bish, M.², Bradley, C.A.³, Duffeck, M.R.⁴, Faske, T.R.⁵, Grabau, Z.⁶, Isakeit, T.⁷, Kemerait, R.C.⁸, Koehler, A.⁹, Langston, D.¹⁰, Lux, L.¹¹, Mueller, J.D.¹², Padgett, G.B.¹³, Plumblee, M.¹², Price, P.P.¹³, Sikora, E.J.¹⁴, Small, I.M.¹⁵, Spurlock, T.⁵, Watson, T.¹³, Young, H.¹⁶, and Zeng, Y.¹⁷

¹Mississippi State University, Stoneville, MS; ²University of Missouri, Columbia, MO;

³University of Kentucky, Princeton, KY; ⁴Oklahoma State University, Stillwater, OK;

⁵University of Arkansas, Lonoke, AR; ⁶University of Florida, Gainesville, FL; ⁷Texas A&M University, College Station, TX; ⁸University of Georgia, Tifton, GA; ⁹University of Delaware, Newark, DE; ¹⁰Virginia Tech, Suffolk, VA; ¹¹North Carolina State University, Raleigh, NC;

¹²Clemson University, Blackville, SC; ¹³Louisiana State University, Baton Rouge, LA; ¹⁴Auburn University, Auburn, AL; ¹⁵University of Florida, Quincy, FL; ¹⁶University of Tennessee, Jackson, TN; ¹⁷Virginia Tech University, Blacksburg, VA

The Southern Soybean Disease Workers (SSDW) have published soybean disease loss estimates for the southern United States since 1974. Summaries of the results from between 1977 and 2019 have been published in numerous refereed scientific journals (7; 11; 16; 18-27; 29-30). The annual losses from between 2015 and 2022 have been presented annually in the SSDW proceedings (1-6; 8-10) and most recently in a publication that included the estimates from 2015 to 2019 in Plant Health Progress that includes the loss estimates from the entire soybean production region including the southern and northern states for a total of 29 states and Ontario, Canada (11). A website through the University of Illinois Extension Service summarizes the estimated yield losses from both the northern and southern U.S. and includes data from 1996 through 2014. The website can be accessed at:

http://extension.cropsci.illinois.edu/fieldcrops/diseases/yield_reductions.php

The additional supporting presentation of loss estimates were included in the annual proceedings of the SSDW as well as some university-related sources (12-15; 17; 28). One last new output source has been at the Crop Protection Network where a soybean disease loss calculator uses data from 1996 through 2021 to provide information related to the percent losses, total bushels lost to disease and specific diseases, loss in dollars, and the potential losses on a per acre basis. The disease loss calculator can be access at:

<https://loss.cropprotectionnetwork.org/crops/soybean-diseases>

The disease loss estimates for the 2023 season contained in the current proceedings document were obtained from representatives across the southern U.S. through various methods. Plant pathologists with soybean pathology responsibilities were queried to provide the estimates of loss from their respective states in November 2023. Most individuals relied on multiple methods to derive estimates. Methods to derive losses included: Field surveys, plant disease diagnostic clinic samples, variety trials, questionnaires to Cooperative Extension staff, research plots, grower demonstrations, private crop consultant reports, foliar fungicide trials, sentinel plot data, variety trial ratings, and "pure guess". To complete the loss estimates for each state, USDA/NASS

production figures were downloaded in January 2024 and production losses were calculated based on estimates of yield in the absence of disease. One additional topic that has recently been included to the presentation of the loss estimates (2018 through 2023) has been a general environmental comparison from each state. However, one major change was made for the 2022 season in that all of the weather station locations were changed to a location collecting weather data from each state's greatest soybean producing county/parish. In situations where the greatest county/parish did not contain a weather station that recorded data the next greatest county/parish was used. Soybean county data were gathered from the USDA Farm Service Agency (FSA) website (<https://www.fsa.usda.gov/news-room/efoia/electronic-reading-room/frequently-requested-information/crop-acreage-data/index>). The data set from 2023 was parsed to determine the greatest counties in each state and irrigated and non-irrigated acres were added up to arrive at those totals. Environmental data from each county were then collected from the National Centers for Environmental Information (<https://www.ncdc.noaa.gov/cdo-web/>) by conducting searches within each state and downloading the entire data set for temperature and precipitation from each corresponding location. State, county/parish, total number of acres within each county/parish, and designated weather station name are presented in Table 1. Environmental data representing the most current 30-year normal (1981-2010) were downloaded for each corresponding location from the National Centers for Environmental Information data tools which includes climate normal (<https://www.ncdc.noaa.gov/cdo-web/datatools/normal>).

Production losses associated with disease severity estimates were based on the formula used to derive production losses:

Potential production without disease loss = actual production ÷ (1-percent loss) (decimal fraction)

Rounding errors may occur in the tables provided below due to the presence of low levels of disease estimated by the state pathologist. Total losses in the form of percent disease loss by state and total losses in millions of bushels were determined by averaging the loss by state with the inclusion of the trace estimates.

The 2023 total acres harvested, average yield (bushels/Acre), and total production (yield in bushels) from each state are presented in Table 2. Soybean acreage in the 16 southern states decreased in 2023 when compared to the 2022 acreage reported by 7% (1). In general, 14 of the southern states reported a decrease in the harvested acres, while two states reported increases. The 2023 average per acre soybean yield was 43.3 bu/A, which was a 0.04% increase compared to the average yield from 2022. Similar to 2022 when several states reported a statewide record yield, none of the states reported a statewide yield in 2023. Even though harvested acres were down across the region, 11 of the states reported increases in the total harvested yield from their respective state between 2023 and 2022. In addition, 15 of the states reported increases in the average state yield, with only Virginia reporting a reduction in average yield. In 2023, more than 925 million bushels were harvested from approximately 19.2 million acres from the 16 southern states accounting for a 4% decrease in the total harvest compared to 2022.

Percentage loss estimates from each state are specific as to causal organism or the common name of the disease (Table 3). The total estimated average percent disease loss for 2023 was lower than

the estimated loss observed during 2022 by 0.9%. As a whole, five states reported an increase in percent disease losses compared to 2022 (AL, KY, MO, NC and TN). The remainder of the 16 states, 11, reported a reduction in percent loss estimates ranging from 0.08% (in OK and VA) to 7.28% (in LA). In terms of the top five diseases encountered during 2023, root-knot nematode, soybean cyst nematode (SCN), charcoal rot, *Cercospora* leaf blight, and the reniform nematode were the top five diseases in order of importance. Three of the top five diseases were similar between 2022 and 2023, with charcoal rot and the reniform nematode being greater issues in 2023 than in 2022 likely as a result of the environment encountered during the season. Breaking the diseases evaluated into categories based on specific plant parts impacted by the diseases within the survey resulted in four categories: Nematode diseases (60%), stem and root diseases (22%), foliar diseases (15%), seedling diseases (5%), and seed diseases (2%). Breaking the diseases down into categories of plant parts impacted helps highlight the importance of specific groups of diseases and which disease areas are causing the greatest estimated losses in a given year/season. Diseases included in the category “other diseases” could not be separated into separate categories and therefore were not included in any single category.

In terms of the disease losses in millions of bushels, the 2023 disease losses accounted for an estimated 49.41, a 15.9% decrease compared to the estimated losses incurred during 2022 (Table 4).

Environmental conditions during 2023 were extremely dry and somewhat warmer than normal across the southern region during June and July (Table 5). In general, a greater number of states recorded reductions in total rainfall over the course of 2023 when compared to the 30-year norm. In all, 10 states recorded negative rainfall totals for 2023 with rainfall totals being more than 74 inches below normal across the region. The decrease accounted for an 9.4% decrease in total rainfall across the region when compared with the 30-year norm. Total rainfall varied greatly by state with 11 states (AL, FL, LA, MD, MO, MS, NC, OK, TN, TX, and VA) recording overall reductions during 2023 when compared with the 30-year normal by between -4.7 inches (TN) and -13.7 inches (NC). The remainder of the states, AR, FL, GA, KY, and SC recorded increases in rainfall compared to the 30-year norm that ranged from 1.2 (SC) to 6.3 (AR). In addition, temperature for 2023 was also compared to the 30-year normal (1981-2010) at each of the locations. In general, looking across the entire year, based on temperature averages for the whole year, three months, May, June, and October were below the 30-year normal temperatures across the region. Conversely, the remainder of the months had temperatures above normal with the greatest temperature increases in January (6.2°F) and February (6.1°F). Looking at temperature data by month, eight months had average temperature increases.

Acknowledgments

Funding was provided from the United Soybean Board through the Crop Protection Network to collate the disease loss estimates from the southern region as part of a larger effort to collect losses as they relate to plant diseases from the entire soybean producing area in the U.S. The members of the SSDW Disease Loss Estimate Committee see value in continuing to collect the estimates on an annual basis and will continue to seek funding to support the effort in the future.

Literature Cited

1. Allen, T. W., Bissonnette, K., Bradley, C. A., Faske, T. R., Grabau, Z., Isakeit, T., Kemerait, R. C., Koehler, A., Langston, D., Lofton, J., Mueller, J. D., Padgett, B. B., Price, P. P., Sikora, E. J., Small, I. M., Vann, R., and Young, H. 2023. Southern United States soybean disease loss estimates for 2022. Pages 1-11 in: Proceedings of the Southern Soybean Disease Workers, Pensacola Beach, FL, March 1-2, 2023.
2. Allen, T. W., Bissonnette, K., Bradley, C. A., Damicone, J. P., Dufault, N. S., Faske, T. R., Isakeit, T., Kemerait, R. C., Koehler, A., Langston, D., Lofton, J., Mueller, J. D., Padgett, G. B., Price, P. P., Sikora, E. J., Small, I. M., Vann, R., and Young, H. 2022. Southern United States soybean disease loss estimates for 2021. Proceedings of the Southern Soybean Disease Workers, Pensacola Beach, FL, March 4-5, 2022.
3. Allen, T. W., Bissonnette, K., Bradley, C. A., Damicone, J. P., Dufault, N. S., Faske, T. R., Isakeit, T., Kemerait, R. C., Koehler, A., Mehl, H. L., Mueller, J. D., Padgett, G. B., Price, P. P., Sikora, E. J., Small, I. M., Thiessen, L., and Young, H. 2021. Southern United States soybean disease loss estimates for 2020. Proceedings of the Southern Soybean Disease Workers, Pensacola Beach, FL, March 4-5, 2021.
4. Allen, T. W., Bissonnette, K., Bradley, C. A., Damicone, J. P., Dufault, N. S., Faske, T. R., Isakeit, T., Kemerait, R. C., Koehler, A., Mehl, H. L., Mueller, J. D., Padgett, G. B., Price, P. P., Sikora, E. J., Small, I. M., Thiessen, L., and Young, H. 2020. Southern United States soybean disease loss estimates for 2019. Proceedings of the Southern Soybean Disease Workers, Pensacola Beach, FL, March 4-5, 2020.
5. Allen, T. W., Bissonnette, K., Bradley, C. A., Damicone, J. P., Dufault, N. S., Faske, T. R., Hollier, C. A., Isakeit, T., Kemerait, R. C., Kleczewski, N. M., Mehl, H. L., Mueller, J. D., Overstreet, C., Price, P. P., Sikora, E. J., Spurlock, T. N., Thiessen, L., and Young, H. 2019. Southern United States soybean disease loss estimates for 2018. Proceedings of the Southern Soybean Disease Workers, Pensacola Beach, FL, March 7-8, 2019.
6. Allen, T. W., Bissonnette, K., Bradley, C. A., Damicone, J. P., Dufault, N. S., Faske, T. R., Hollier, C. A., Isakeit, T., Kemerait, R. C., Kleczewski, N. M., Mehl, H. L., Mueller, J. D., Overstreet, C., Price, P. P., Sikora, E. J., Spurlock, T. N., Thiessen, L., and Young, H. 2018. Southern United States soybean disease loss estimates for 2017. Proceedings of the Southern Soybean Disease Workers, Pensacola Beach, FL, March 7-8, 2018.
7. Allen, T. W., Bradley, C. A., Sisson, A. J., Byamukama, E., Chilvers, M. I., Coker, C. M., Collins, A. A., Damicone, J. P., Dorrance, A. E., Dufault, N. S., Esker, P. D., Faske, T. R., Giesler, L. J., Grybauskas, A. P., Hershman, D. E., Hollier, C. A., Isakeit, T., Jardine, D. J., Kemerait, R. C., Kleczewski, N. M., Koenning, S. R., Kurle, J. E., Malvick, D. K., Markell, S. G., Mehl, H. L., Mueller, D. S., Mueller, J. D., Mulrooney, R. P., Nelson, B. D., Newman, M. A., Osborne, L., Overstreet, C., Padgett, G. B., Phipps, P. M., Price, P. P., Sikora, E. J., Smith, D. L., Spurlock, T. N., Tande, C. A., Tenuta, A. U., Wise, K. A., Wrather, J. A., and Young-Kelly, H. 2017. Soybean yield loss estimates due to diseases in the United States and Ontario, Canada from 2010 to 2014. *Plant Health Progress*. doi: 10.1094/PHP-RS-16-0066.
8. Allen, T. W., Bradley, C. A., Damicone, J. P., Dufault, N. S., Faske, T. R., Hollier, C. A., Isakeit, T., Kemerait, R. C., Kleczewski, N. M., Kratochvil, R. J., Mehl, H. L., Mueller, J. D., Overstreet, C., Price, P. P., Sikora, E. J., Spurlock, T. N., Thiessen, L., Wiebold, W. J.,

- and Young, H. 2017. Southern United States soybean disease loss estimates for 2016. Proceedings of the Southern Soybean Disease Workers, Pensacola Beach, FL, March 8-9, 2017.
9. Allen, T.W., Damicone, J. P., Dufault, N. S., Faske, T. R., Hershman, D. E., Hollier, C. A., Isakeit, T., Kemerait, R. C., Kleczewski, N. M., Koenning, S. R., Mehl, H. L., Mueller, J. D., Overstreet, C., Price, P. P., Sikora, E. J., and Young, H. 2016. Southern United States soybean disease loss estimate for 2015. Pages 11-15 in: Proceedings of the Southern Soybean Disease Workers, Pensacola Beach, FL, March 9-10, 2016.
 10. Allen, T.W., Damicone, J. P., Dufault, N. S., Faske, T. R., Hershman, D. E., Hollier, C. A., Isakeit, T., Kemerait, R. C., Kleczewski, N. M., Koenning, S. R., Mehl, H. L., Mueller, J. D., Overstreet, C., Price, P. P., Sikora, E. J., and Young, H. 2015. Southern United States soybean disease loss estimate for 2014. Pages 10-15 in: Proceedings of the Southern Soybean Disease Workers, Pensacola Beach, FL, March 11-12, 2015.
 11. Bradley, C. A., Allen, T. W., Sisson, A. J., Bergstrom, G. C., Bissonnette, K. M., Bond, J., Byamukama, E., Chilvers, M. I., Collins, A. A., Damicone, J. P., Dorrance, A. E., Dufault, N. S., Esker, P. D., Faske, T. R., Fiorellino, N. M., Giesler, L. J., Hartman, G. L., Hollier, C. A., Isakeit, T., Jackson-Ziems, T. A., Jardine, D. J., Kelly, H. M., Kemerait, R. C., Kleczewski, N. M., Koehler, A. M., Kratochvil, R. J., Kurle, J. E., Malvick, D. K., Markell, S. G., Mathew, F. M., Mehl, H. L., Hehl, K. M., Mueller, D. S., Mueller, J. D., Nelson, B. D., Overstreet, C., Padgett, G. B., Price, P. P., Sikora, E. J., Small, I., Smith, D. L., Spurlock, T. N., Tande, C. A., Telenko, D. E. P., Tenuta, A. U., Thiessen, L. D., Warner, F., Wiebold, W. J., and Wise, K. A. 2021. Soybean yield loss estimates due to diseases in the United States and Ontario, Canada, from 2015 to 2019. *Plant Health Prog.* 22:483-495.
 12. Koenning, S. R. 2014. Southern United States soybean disease loss estimate for 2013. Pages 1-5 in: Proceedings of the Southern Soybean Disease Workers, Forty-First Annual Meeting, Pensacola, FL.
 13. Koenning, S. R. 2013. Southern United States soybean disease loss estimate for 2012. Pages 1-5 in: Proceedings of the Southern Soybean Disease Workers, Fortieth Annual Meeting, Pensacola, FL.
 14. Koenning, S. R. 2012. Southern United States soybean disease loss estimate for 2011. Pages 1-5 in: Proceedings of the Southern Soybean Disease Workers, Thirty-Ninth Annual Meeting, Pensacola, FL.
 15. Koenning, S. R. 2011. Southern United States soybean disease loss estimate for 2010. Pages 1-5 in: Proceedings of the Southern Soybean Disease Workers, Thirty-Eighth Annual Meeting, Pensacola, FL.
 16. Koenning, S. R., and Wrather, J. A. 2010. Suppression of soybean yield potential in the Continental United States by plant diseases from 2006 to 2009. *Plant Health Progress* doi:10.1094-2010-1122-01-RS.
 17. Koenning, S. R. 2010. Southern United States soybean disease loss estimate for 2009. Pages 1-5 in: Proceedings of the Southern Soybean Disease Workers, Thirty-Seventh Annual Meeting, Pensacola, FL.
 18. Mulrooney, R. P. 1988. Soybean disease loss estimate for Southern United States in 1985 and 1986. *Plant Disease* 72:364-365.
 19. Mulrooney, R. P. 1988. Soybean disease loss estimate for Southern United States in 1987. *Plant Disease* 72:915.

20. Pratt, P. W., and Wrather, J. A. 1998. Soybean disease loss estimates for the Southern United States during 1994-1996. *Plant Disease* 82:114-116.
21. Sciumbato, G. L. 1993. Soybean disease loss estimate for the Southern United States during 1988-1991. *Plant Disease* 77:954-956.
22. Whitney, G. 1978. Southern states soybean disease loss estimates 1977. *Plant Disease Reporter* 62:1078-1079.
23. Wrather, J. A., Chambers, A. Y., Fox, J. A., Moore, W. F., and Sciumbato, G. L. 1995. Soybean disease loss estimates for the southern United States, 1974 to 1994. *Plant Disease* 79:1076-1079.
24. Wrather, J. A., and Sciumbato, G. L. 1995. Soybean disease loss estimates for the Southern United States during 1992-1993. *Plant Disease* 79:84-85.
25. Wrather, J. A., Anderson, T. R., Arsyad, D. M., Gai, J., Ploper, L. D., Porta-Puglia, A., Ram, H. H., and Yorinori, J. T. 1997. Soybean disease loss estimates for the top 10 soybean producing countries in 1994. *Plant Disease* 81:107-110.
26. Wrather, J. A., Stienstra, W. C., and Koenning, S. R. 2001. Soybean disease loss estimates for the United States from 1996-1998. *Canadian Journal of Plant Pathology* 23:122-131.
27. Wrather, J. A., Koenning, S. R., and Anderson, T. 2003. Effect of diseases on soybean yields in the United States and Ontario 1999-2002. Online. *Plant Health Progress* doi: 10.1094/PHP-2003-0325-01-RV.
28. Wrather, J. A., and Koenning, S. R. 2005. Soybean disease loss estimates for the United States 1996-2004. <http://aes.missouri.edu/delta/research/soyloss.stm>
29. Wrather, J. A., and Koenning, S. R. 2006. Estimates of disease effects on soybean yields in the United States 2003-2005. *Journal of Nematology* 38:173-180.
30. Wrather, J. A., and Koenning, S. R. 2009. Effects of diseases on soybean yields in the United States 1996 to 2007. *Plant Health Progress* doi:10.1094/PHP-2009-0401-01-RS.

Table 1. Locations where environmental data were downloaded based on the county/parish in each state with the greatest soybean production and a town within that county that contained environmental data for the 2023.

2023 Acres of soybean production ^b				
State	County/Parish ^a	Location	soybean production ^b	Specific weather station ^c
Alabama	Limestone	Athens	49,966	Athens
Arkansas	Mississippi	Keiser	241,824	Keiser
Delaware	Sussex	Georgetown	70,987	Georgetown Delaware Coastal Airport
Florida	Jackson	Marianna	3,660	Marianna Municipal Airport
Georgia	Laurens	Dublin	9,375	Dublin 2
Kentucky	Christian	Hopkinsville	85,712	Hopkinsville (Christian County)
Louisiana	Madison	Tallulah	82,180	Tallulah Vicksburg Regional Airport
Maryland	Worcester	Snow Hill	4,099	Baltimore Washington International Airport
Mississippi	Bolivar	Cleveland	276,311	Cleveland
Missouri	Saline	Marshall	142,310	Marshall
North Carolina	Robeson	Lumberton	80,465	Lumberton
Oklahoma	Kay	Blackwell	90,131	Blackwell 4 SSE Mesonet
South Carolina	Florence	Florence	40,882	Florence Regional Airport
Tennessee	Gibson	Milan	102757	Milan Experimental Station
Texas	Lamar	Paris	16,082	Paris
Virginia	Hanover	Hanover	21,180	Ashland Hanover Co. Municipal Airport

^a County or parish soybean production data from each of the greatest soybean producing counties/parishes were determined based on data provided from the crop acreage data set from the USDA Farm Service Agency (www.fsa.usda.gov). Note that a weather station could not be located in the greatest soybean producing county/parish in Louisiana, Maryland, Missouri, or Virginia. The third largest soybean producing parish in Louisiana, 18th largest soybean producing county in Maryland, and the third largest in Missouri and Virginia were therefore relied on for environmental data.

^b The total soybean acres produced in each county were determined based on FSA data and were added in instances where soybean production was separated based on irrigation practice, seed production, or the production of edamame.

^c Specific weather station names are included for the purposes of presenting a historical record of these data as downloaded from the National Centers for Environmental Information website (www.ncdc.noaa.gov). In addition, weather stations were chosen that included a 30-year normal data set so comparisons could be made between 2022 and the normal environmental data.

The specific weather station was based on the name assigned by NASS and can be located on their website.

^d The data for the location in Kentucky was downloaded from http://kymesonet.org/monthly_summaries.html. The Kentucky mesonet site was the only one that included temperature as well as precipitation data.

Table 2. Soybean production from the 16 southern states during 2023 ^a.

State	Acres (1,000s) ^b	Bu/Acre ^{b,c}	Yield in bu/A (1,000s) ^b
Alabama	345 (-)	43 (+2)	14,835 (+)
Arkansas	2,950 (-)	54* (+2)	159,300 (-)
Delaware	148 (-)	46 (+2)	6,808 (+)
Florida	18 (-)	43 (-9)	774 (-)
Georgia	155 (-)	43* (+2)	6,665 (+)
Kentucky	1,820 (-)	55 (+4)	100,100 (+)
Louisiana	980 (-)	40 (-7)	39,200 (-)
Maryland	460 (-)	47 (+4)	21,620 (-)
Mississippi	2,130 (-)	56 (+2)	119,280 (-)
Missouri	5,520 (-)	48 (+2.5)	264,960 (-)
North Carolina	1,630 (-)	38.5 (-)	62,755 (-)
Oklahoma	410 (+)	26 (+9)	10,660 (+)
South Carolina	385 (-)	39* (+2)	15,015 (+)
Tennessee	1,570 (-)	51* (+3)	80,070 (+)
Texas	85 (-)	25 (+5)	2,125 (+)
Virginia	570 (-)	38 (-3)	21,660 (-)
TOTAL	19,176 (-)		925,827 (-)
Avg. 43.3 (+1.9)			

^a Data were compiled from the USDA National Agricultural Statistics Service Crop Production 2023 Summary as distributed in January 2024. The report, in its entirety, can be downloaded from: <https://downloads.usda.library.cornell.edu/usda-esmis/files/k3569432s/ns065v292/8910md644/cropan24.pdf>.

Additional data were supplemented for states such as Florida by using the FSA acres information which can be accessed at: <https://www.fsa.usda.gov/newsroom/efoia/electronic-reading-room/frequently-requested-information/crop-acreage-data/index>

^b Difference from 2022 indicated in parentheses as either a decrease (-) or increase (+).

Table 3. Estimated percentage loss of soybean yield due to diseases from 16 southern states during 2023.

Disease	% yield suppression by state																AVG
	AL ^a	AR	DE	FL	GA	KY	LA	MD	MO	MS	NC	OK	SC	TN	TX	VA	
Anthracnose	0.00	0.00	0.00	0.01	0.10	0.00	0.00	0.00	0.00	0.00	0.01	0.00	0.01	0.40	0.00	0.01	0.03
Bacterial diseases	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.01	0.00	0.00	0.00	0.00
Brown stem rot	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.01	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Cercospora leaf blight	1.00	0.25	0.01	0.10	1.00	0.10	0.50	0.01	0.05	1.00	0.75	0.10	0.50	0.01	0.00	0.25	0.35
Charcoal rot	1.00	0.01	0.00	0.01	0.01	0.02	0.00	0.00	0.60	0.00	0.05	7.00	0.01	0.50	0.00	0.01	0.58
Downy mildew	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.01	0.00	0.00	0.00	0.00	0.01	0.00
Frogeye leaf spot	0.01	0.03	0.00	0.00	0.00	1.00	0.02	0.00	0.00	0.00	0.08	0.00	0.01	1.40	0.00	0.03	0.16
Fusarium wilt and root rot	0.00	0.01	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.05	0.00	0.00	0.00	0.00	0.01	0.00
Other diseases ^b	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Phomopsis seed decay	0.02	0.00	0.01	0.10	0.10	0.00	0.00	0.01	0.00	0.01	0.00	0.01	0.50	0.01	0.00	0.10	0.05
Phytophthora root and stem rot	0.00	0.02	0.00	0.00	0.00	0.60	0.00	0.00	0.60	0.00	0.15	0.00	0.00	0.05	0.00	0.00	0.09
Pod and stem blight	0.01	0.01	0.00	0.10	0.50	0.05	0.00	0.00	0.10	0.00	0.10	0.00	0.50	0.05	0.00	0.01	0.09
Purple seed stain	0.02	0.05	0.05	0.10	0.10	0.03	0.01	0.05	0.00	0.02	0.01	0.10	0.01	0.08	0.00	0.01	0.04
Red crown rot	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.10	0.00	0.00	0.00	0.01
Reniform nematode	0.30	0.30	0.00	0.50	0.10	0.00	1.90	0.00	0.00	0.06	0.00	0.00	0.50	0.01	0.00	0.00	0.23
Root-knot nematode	1.50	4.30	1.50	3.00	1.50	0.00	2.25	0.50	0.01	1.85	2.00	0.00	2.50	0.01	0.00	1.00	1.37
Soybean cyst nematode	0.05	0.30	2.50	0.00	0.00	2.80	0.00	1.00	2.50	0.05	2.00	0.00	0.50	1.80	0.00	2.00	0.97
Other nematodes ^c	0.00	0.03	0.05	0.10	0.10	0.00	0.00	0.01	0.00	0.00	0.00	0.00	2.50	0.00	0.00	0.25	0.19
Rhizoctonia aerial blight	0.01	0.08	0.00	0.00	0.00	0.00	0.10	0.00	0.00	0.10	0.00	0.00	0.00	0.00	0.00	0.00	0.02
Sclerotinia stem rot (white mold - <i>Sclerotinia sclerotiorum</i>)	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Seedling diseases	0.00	0.03	0.01	0.00	0.00	0.70	0.00	0.01	0.01	0.30	0.30	0.00	0.02	1.00	0.00	0.25	0.16
Septoria brown spot	0.00	0.02	0.02	0.00	0.00	0.40	0.02	0.02	0.00	0.10	0.05	0.00	0.05	1.00	0.00	0.05	0.11
Southern blight	0.05	0.01	0.00	0.01	0.01	0.00	0.03	0.00	0.00	0.30	0.02	0.00	0.05	0.00	0.00	0.00	0.03
Soybean rust	0.01	0.00	0.00	0.01	0.50	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.03
Stem canker	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.20	0.10	0.00	0.00	0.00	0.02	0.02
Sudden death syndrome	0.00	0.00	0.00	0.00	0.00	1.00	0.00	0.00	0.30	0.00	0.01	0.00	0.00	0.01	0.00	0.02	0.08
Taproot decline	0.05	0.02	0.00	0.00	0.00	0.00	0.50	0.00	0.00	0.30	0.00	0.00	0.01	0.00	0.00	0.00	0.06
Target spot	0.00	0.00	0.00	0.00	0.00	0.10	0.00	0.00	0.00	0.00	0.01	0.00	0.15	0.15	0.00	0.01	0.03
Virus Diseases ^d	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.01	0.00	0.04	0.00	0.00	0.00	0.00
Total disease %	4.03	5.47	4.14	4.04	4.02	6.80	5.33	1.60	4.18	4.09	5.81	7.32	7.97	6.46	0.00	4.04	4.71

^a Rounding errors may exist since some numbers presented carry decimal places beyond the hundredths place.^b Other diseases listed included: Phymatotrichopsis root rot (TX).^c Other nematodes listed included: Columbia lance nematode (SC), lance nematode (AR, DE, MD), lesion nematode (DE, MD), sting nematode (GA), stubby root nematode (SC).^d Virus diseases listed included: *Bean pod mottle virus* (KY), *Soybean vein necrosis virus* (DE, KY, MD, MS), *Tobacco ringspot virus* (KY).

Table 4. Estimated suppression of soybean yield (Millions of Bushels) as a result of disease during 2023.

Disease	yield suppression by state (millions of bushels) ^a																TOTAL
	AL	AR	DE	FL	GA	KY	LA	MD	MS	MO	NC	OK	SC	TN	TX	VA	
Anthracnose	0.00	0.00	0.00	0.00	0.01	0.00	0.00	0.00	0.00	0.00	0.01	0.00	0.00	0.34	0.00	0.00	0.36
Bacterial diseases	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Brown stem rot	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.03	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.03
Cercospora leaf blight	0.15	0.42	0.00	0.00	0.07	0.11	0.21	0.00	0.14	1.24	0.50	0.01	0.08	0.00	0.00	0.06	3.00
Charcoal rot	0.15	0.02	0.00	0.00	0.00	0.02	0.00	0.00	1.66	0.00	0.03	0.81	0.00	0.43	0.00	0.00	3.12
Downy mildew	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.01	0.00	0.00	0.00	0.00	0.00	0.01
Frogeye leaf spot	0.00	0.05	0.00	0.00	0.00	1.07	0.01	0.00	0.00	0.00	0.05	0.00	0.00	1.20	0.00	0.01	2.39
Fusarium wilt and root rot	0.00	0.01	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.03	0.00	0.00	0.00	0.00	0.00	0.05
Other diseases	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Phomopsis seed decay	0.00	0.00	0.00	0.00	0.01	0.00	0.00	0.00	0.00	0.01	0.00	0.00	0.08	0.00	0.00	0.02	0.14
Phytophthora root and stem rot	0.00	0.03	0.00	0.00	0.00	0.64	0.00	0.00	1.66	0.00	0.10	0.00	0.00	0.04	0.00	0.00	2.48
Pod and stem blight	0.00	0.02	0.00	0.00	0.03	0.05	0.00	0.00	0.28	0.00	0.07	0.00	0.08	0.04	0.00	0.00	0.58
Purple seed stain	0.00	0.08	0.00	0.00	0.01	0.03	0.00	0.01	0.00	0.02	0.01	0.01	0.00	0.06	0.00	0.00	0.26
Red crown rot	0.00	0.00	0.00	0.00	0.00	0.00	0.79	0.00	0.00	0.00	0.00	0.00	0.02	0.00	0.00	0.00	0.80
Reniform nematode	0.05	0.51	0.00	0.00	0.01	0.00	0.93	0.11	0.00	0.07	0.00	0.00	0.08	0.01	0.00	0.00	1.76
Root-knot nematode	0.23	7.25	0.11	0.02	0.10	0.00	0.00	0.22	0.03	2.30	1.33	0.00	0.41	0.01	0.00	0.23	12.24
Soybean cyst nematode	0.01	0.51	0.18	0.00	0.00	3.01	0.00	0.00	6.91	0.06	1.33	0.00	0.08	1.54	0.00	0.45	14.08
Other nematodes ^b	0.00	0.05	0.00	0.00	0.01	0.00	0.04	0.00	0.00	0.00	0.00	0.00	0.41	0.00	0.00	0.06	0.57
Rhizoctonia aerial blight	0.00	0.13	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.12	0.00	0.00	0.00	0.00	0.00	0.00	0.26
Sclerotinia stem rot (white mold - <i>Sclerotinia sclerotiorum</i>)	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Seedling diseases	0.00	0.05	0.00	0.00	0.00	0.75	0.01	0.00	0.01	0.37	0.20	0.00	0.00	0.86	0.00	0.06	2.32
Septoria brown spot	0.00	0.03	0.00	0.00	0.00	0.43	0.01	0.00	0.00	0.12	0.03	0.00	0.01	0.86	0.00	0.01	1.51
Southern blight	0.01	0.02	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.37	0.01	0.00	0.01	0.00	0.00	0.00	0.42
Soybean rust	0.00	0.00	0.00	0.00	0.03	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.04
Stem canker	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.13	0.01	0.00	0.00	0.00	0.00	0.15
Sudden death syndrome	0.00	0.00	0.00	0.00	0.00	1.07	0.21	0.00	0.83	0.00	0.01	0.00	0.00	0.01	0.00	0.00	2.13
Taproot decline	0.01	0.03	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.37	0.00	0.00	0.00	0.00	0.00	0.00	0.42
Target spot	0.00	0.01	0.00	0.00	0.00	0.11	0.00	0.00	0.00	0.00	0.01	0.00	0.02	0.13	0.00	0.00	0.28
Virus Diseases ^c	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.01	0.00	0.01	0.00	0.00	0.00	0.01
Total disease %	0.62	9.22	0.29	0.03	0.28	7.31	2.21	0.35	11.56	5.08	3.87	0.84	1.30	5.53	0.00	0.91	49.41

^aRounding errors may exist since some numbers presented carry decimal places beyond the hundredths place.^cOther nematodes listed included: Columbia lance nematode (SC), lance nematode (DE), lesion nematode (AL, AR, DE, KY, SC), spiral (KY, LA), sting nematode (GA, SC), stubby root nematode (SC).^cVirus diseases listed included: *Soybean vein necrosis virus* (DE, KY, MD, MS, NC).

Table 5. Deviation of the 2023 temperature from the 30-year normal and the total precipitation for 2023 and the 30-year normal from each of the 16 southern soybean producing states based on data downloaded from the centroid for each respective state.

State	Deviation from the 30-year temperature norm (°F) ^a												Total precip (in) ^b		
	January	February	March	April	May	June	July	August	September	October	November	December	2023	30-year	Deviation
Alabama	3.5	3.9	-2.3	-3.4	-1.7	-1.1	-0.4	-1.0	1.4	-1.2	-0.1	1.2	52.1 (5)	59.1	-7.0
Arkansas	6.3	3.3	-1.4	-1.1	0.0	-1.5	-2.0	-2.4	0.8	1.6	0.9	4.2	58.5 (5)	52.2	6.3
Delaware	9.7	8.5	3.5	6.1	-1.4	-0.6	0.2	-1.0	-0.1	2.4	-0.7	3.3	45.7 (4)	43.8	1.8
Florida	6.0	7.2	2.0	-0.5	-2.4	-0.9	0.4	5.5	0.3	0.0	-1.2	-0.5	45.6 (4)	51.1	-5.5
Georgia	5.1	9.0	2.3	1.2	-4.1	-2.4	-0.7	1.9	-1.6	-0.3	-1.2	1.3	52.8 (6)	49.3	3.5
Kentucky	7.9	8.8	0.7	0.2	-0.3	-1.4	-1.9	-3.6	-0.7	-1.2	2.5	5.1	54.7 (7)	51.8	3.0
Louisiana	6.9	6.4	4.3	-1.6	1.5	1.7	1.6	7.5	4.9	3.0	0.5	4.6	44.2 (2)	53.3	-9.2
Maryland	3.7	4.0	-1.1	2.7	-2.9	-1.7	1.0	-0.5	-0.8	0.2	-2.7	-0.2	42.3 (5)	47.4	-5.1
Missouri	7.9	6.3	-1.3	5.3	4.2	3.2	2.3	0.4	4.6	4.1	3.4	8.8	34.5 (6)	41.8	-7.3
Mississippi	7.1	4.2	0.7	-1.8	-0.5	-0.5	-0.8	1.2	-7.2	-11.7	-3.6	7.8	50.2 (4)	59.3	-9.1
North Carolina	6.3	7.7	2.0	1.1	-4.1	-3.4	1.0	2.0	-0.5	0.6	-0.7	3.7	37.2 (3)	50.8	-13.7
Oklahoma	4.3	3.1	-1.1	1.7	0.4	1.2	0.3	3.0	5.6	0.1	2.7	.	26.8 (2)	38.8	-12.1
South Carolina	5.3	6.7	1.1	0.8	-4.0	-3.0	0.1	1.7	-1.0	0.2	-0.3	2.6	46.5 (7)	45.3	1.2
Tennessee	6.1	5.3	0.2	-1.9	-0.7	-0.6	-1.8	-3.5	-0.1	-0.2	1.0	6.1	51.5 (5)	56.2	-4.7
Texas	6.3	3.3	1.7	-2.4	-0.1	-2.1	-2.6	3.5	2.2	-1.8	1.2	3.5	43.8 (5)	48.9	-5.1
Virginia	6.7	9.1	4.1	66.6	-0.6	-0.7	2.9	3.3	1.9	3.1	0.9	3.8	35.4 (3)	47.2	-11.8
Avg.	6.2	6.1	1.0	4.6	-1.0	-0.9	0.0	1.1	0.6	-0.1	0.2	3.7	--	--	-74.8

^a Deviations of temperature were calculated based on subtracting the average temperature for each month from the 30-year normal. Negative numbers are deviations below the normal and positive numbers are deviations above the normal temperature for the 30-year period from 1981-2010.

^b Numbers in parentheses equal the number of months where the total rainfall was greater than the 30-year normal for each month at the given location. Only values greater than 0.0 were considered “above the normal”.