

### Crop Replant and Rotation Guide for Soybean Herbicides\*

Herbicide	Replant/Crop Rotation	Time Interval	Precautions
Anthem Maxx	S,C CT,P,SF,W AL,R SG	I 4 months 10 months 11 months	See label for rotational intervals based on rates applied.
Assure II/Targa	S,C All	I 120 days	
Authority MTZ	S B,W,SG R AL,CT,GS,P,SF	I 4 months 10 months 12 months	See label for pH restrictions. High pH = longer rotations.
Authority Edge/ Authority Supreme	S, SF C,P,W GS,R B AL,CT,O	I 4 months 10 months 11 months 12 months	See label for pH, rate restrictions. High pH rates have longer rotational intervals.
Authority XL	B,R,Y,W AL,CL,CT, FC,GS,O, P,R,SC,SF All	4 months 18 months  36 months	See label for pH, rate restrictions.
Basagran	All	I	
Broadaxe XC or Authority Elite	S,SU,DSP P W FC,R,GS O,CT* SC,PC	I 4 months 4.5 months 10 months 12 months 18 months	
Butyrac/Butoxone (2,4-DB)			No restrictions.
Cadet	All	I	
Classic	S SG,RG P C,CT CL,GS,R All	I 3 months 6 months 8 months 9 months 18 months†	If applied after Aug. 1, extend recrop interval for C, CT, R and GS 2 months. † Successful field bioassay must be completed prior to planting. Wait 9 months before trying a bioassay.

Herbicide	Replant/Crop Rotation	Time Interval	Precautions
Cobra			No restrictions.
Dual Magnum or Sequence	S,C,CT,GS† SG,W Rice All	I 4.5 months Next spring 18 months	† Use Concep-treated grain sorghum seed.
Enlist One	S, C CT All	7-14 days 1 month 1 month	Rotation intervals based on rate – see label. Enlist corn, cotton, and soybean can be planted immediately.
Enlist Duo	S, C, CT All	1 month 1 month	Enlist corn, cotton, and soybean can be planted immediately.
Engenia	C GS, S, SG All	I 14-28 days 120 days	Rotation intervals based on rate – see label. Dicamba-tolerant soybean and cotton can be planted immediately.
Enlite	W CT,C,R,SF AL,CL	4 months 9 months 12 months	
Envive	W CT,C,GS,R,AL,CL SF,CA	4 months 10 months 18 months	See label for pH and geographical restrictions. High pH soils have longer rotational intervals.
FirstRate	W C,CT,P,R,GS	3 months 9 months	All others – 30 months plus bioassay.
Flexstar GT 3.5	CT,S W C,P,R,GS Other	0 months 4 months 10 months 18 months	See Reflex.
Fusilade DX or Fusion	S, CT All	I 2 months	
Glufosinate	C,S,CT,R,CA W,B,O GS	I 70 days 180 days	No restrictions.
Glyphosate	All	I	
Intimidator	S B, W C, R CT, RY GS All	I 4.5-8 months 10 months 12 months 18 months 18 months	
Metribuzin	S† AL,C,FG,W,B CT,R All (except root crops)†† Root crops	I 4 months 8 months 12 months 18 months	† Waiting period for replanting soybeans depends on the rate of metribuzin used. See specific label for more information. Add 2 months to time intervals if pH of soil is above 7.5. †† Cover crops may be planted any time, but stand reductions may occur.

\*This table applies to the major field and forage crops. Refer to the herbicide labels for the latest recrop and rotation information for horticultural crops. These are written as best we could interpret the labels. We regret any omissions or errors. Always refer to product labels before using a pesticide or replanting into treated fields.

NOTE: See page 44 for key to abbreviations.

(continued on page 44)

### Crop Replant and Rotation Guide for Soybean Herbicides\* [cont.]

Herbicide	Replant/Crop Rotation	Time Interval	Precautions
Outlook	C,S SG All	I 4 months FY	Do not retreat field with second application.
Paraquat			No restrictions.
Poast/Poast Plus			See label.
Prefix	S CT AL,W,O,R,RY,B GS,R,C	I 1 month 4.5 months 10 months	Use Concep-treated grain sorghum seed. Do not use Reflex or Flexstar in-season.
Prowl	S,CT W,B All	I 4 months† FY	Do not rework soil deeper than treated zone. † Cannot replant using no-tillage practices.
Pursuit	S B,W C Rice All	I† 4 months 9.5 months 18 months 18 months	† Do not rework soil deeper than 2 inches. Do not apply Classic, Lorox Plus, Scepter, Squadron or Tri-Scept the same year as Pursuit or injury to following crops may occur.
Python	AL,P,B,W,RY,O R C GS CT, Sunflower	4 months 6 months 1 month 12 months 18 months	Requires successful bioassay.
Reflex or Flexstar	S SG, W C,CT,R,P,GS All	I 4 months 10 months 18 months	
Resource	SG S,C All	120 days I 30 days	
Reviton	C, W S CT All	I 0 to 7 days 7 to 14 days 120 to 150 days	See label.
Select Max			See label.
Sharpen	C,GS,SG,S,R CT SF, others	I 1.5 months 4 months	† Rotation intervals based on rate – see label.
Sonic/Authority First	W C,R CT,AL,B,O, RY,GS	4 months 10 months 12 months	See label for pH restrictions.
Storm	All	Fall	
Surveil	S W C,P,R,GS SF	I 3 months 9 months 30 days	
Tavium	S CT C B, O, RY, W AL, DSP, GS, P, SF CL R All	28 days 42 days 4 months 4.5 months 6 months 9 months Next spring 12 months	Dicamba-tolerant soybean and cotton can be planted immediately.
Tendovo	S W FC, PC, AL, R B, CT, P, GS SC, All SF	I 4.5 months 9 months 12 months 18 months 30 months	Cover crops for soil building or erosion control may be planted any time but do not graze or harvest for food or feed.

Herbicide	Replant/Crop Rotation	Time Interval	Precautions
Treflan/others	S,CT W,B All	I Fall FY	
Treflan (2X rate)	S,CT R All	I FY 2 years	
Trivence	S W C R CT All	I 4 months 10 months 12 months 18 months 18 months	Rotational intervals based on pH and rates – see label.
Typhoon	W C,CT,R GS and all others	4 months 10 months 18 months	
Ultra Blazer	S All	I Fall	
Valor	S,P W,C,R,GS CT B AL,O,CL,CA	I 30 days 21 days 4 months 8 months	Pre-emerge treatments may injure soybeans.
Verdict	C,S, GS CT,R W	I FY 4 months	† Rotation intervals based on rate – see label.
Warrant	CT,C,GS,S W R, others	I 4 months FY	Do not use more than 3.0 lb/A/year.
Xtendimax	C, S GS CT SG R All	14-28 days 15 days 21 days 22-45 days 120 days 120 days	Rotation intervals based on rate – see label. Dicamba-tolerant soybean and cotton can be planted immediately.
Zidua	C,S W CT GS R, others	I 1 month 2 months 6 months FY	
Zidua Pro	S P, W C AL B, RY CT, GS, SF All	I 4 months 8.5 months 10 months 11 months 18 months 40 months	Following 40 months, before planting any crop not listed on the label, a field bioassay must be completed. See label for Clearfield crop rotation intervals.

#### KEY

##### Crop

All = All crops not specified  
 AL = Alfalfa  
 B = Barley  
 C = Corn  
 CA = Canola  
 CL = Clovers  
 CT = Cotton  
 DSP = Dry Shelled Peas  
 FC = Field Corn

FG = Forage Grasses  
 FL = Forage Legumes  
 GS = Grain Sorghum  
 O = Oats  
 P = Peanuts  
 PC = Popcorn  
 R = Rice  
 RG = Ryegrass  
 RY = Rye

S = Soybean  
 SC = Sweet Corn  
 SF = Sunflower  
 W = Wheat

##### Timing

I = Immediately  
 FY = Following year  
 (usually spring or following fall - 11 to 16 months)

\*This table applies to the major field and forage crops. Refer to the herbicide labels for the latest recrop and rotation information for horticultural crops. These are written as best we could interpret the labels. We regret any omissions or errors. Always refer to product labels before using a pesticide or replanting into treated fields.