

Evaluating New Production Inputs that will contribute to High Yield Soybeans. Project 28-2020

PIs: Southern Ag - Alan Blaine, alan@southernaginc.com; Dan Prevost, dan@southernaginc.com; Christina Hall, christina@southernaginc.com; Zach Reynolds, zach@southernaginc.com; Will McCarty, willm4590@comcast.net; Mitt Wardlaw, mitt@southernaginc.com; Brian Ward, boz@southernaginc.com; W. R. Thompson, bobthompson54@yahoo.com; William Kingery, wlk@msstate.edu.

RATIONALE/JUSTIFICATION FOR RESEARCH:

Much of the current research focuses on specific factors within a cropping system. Many growers have achieved higher than average yields through intensive management. However, many feel they have reached a “glass ceiling” with regards to yield and profit. While there are many acres that can still be improved; it is those that have hit a “glass ceiling” that ask questions regarding how do we break through this barrier. Through observation and discussion with growers, there are a myriad of factors that limit yield. Oftentimes, these factors are things that cannot be adjusted mid-season such as drainage. With that being said, growers in some high yield environments want to increase profitability through different mid-season management strategies.

Several years ago, we observed a soybean crop planted next to cotton. The grower was spraying mepiquat chloride (Pix) plus a plant bug material in the cotton that drifted onto the soybeans. When he began harvest, the soybeans next to the cotton had increased yields and declined as he moved farther from the drifted area. Questions were asked then and in numerous other situations about plant growth regulators and the effect of a cotton spray program on soybeans (materials specifically being applied for plant bugs). In years past we have tried to use Pix on soybeans, but observed no visual results. No one feels plant bugs are impacting soybean growth/ development, but they are considered a major pest in cotton. Based on past observations and numerous questions we would like to evaluate some different insect control strategies combined with growth regulators to see if we can duplicate past observations and improve soybean yields even more.

OBJECTIVE(S):

Soybeans are being grown under management that was unheard of in years past. Although, a lot of inputs go into making a crop, additional inputs need to be evaluated to determine their place in

a production system. In the case of high yields, many growers feel they have maxed out or hit a ceiling. Many produce high yields, but would like to do so consistently. Statewide there are still strides that can be made in soil fertility coupled with some possible micronutrient mixes, but in a lot of situations fertility does not appear to be the limiting factor. For any cropping system, we must first identify the most yield limiting factor. As we identify and solve for these initial limiting factors, other factors will then reveal themselves and the process is repeated. The proposed research will let us look at some strategies that have not been evaluated, but discussed quite a lot over the years.

1. Determine efficacy of higher use rates of Dimilin insecticide in providing season long control of pests.
2. Evaluate the effect of plant bug control on soybean yield using insecticide (Transform) application and timing strategies.
3. Evaluate the effect of plant growth regulators (Pix, Apogee, and Stance) on soybean physiology and yield.
4. Evaluate combinations of treatments listed in Objectives 1-3 for synergism and opportunities to minimize trips across the field.

APPROACH AND EXPERIMENT CONDUCT:

This research is an attempt to identify some new production inputs that will allow growers to achieve higher soybean yields. We implemented three trials: one on a highly fertilized/ leveled field in the hill area and two on high yield potential soil in the MS Delta. Progressive management practices were followed surrounding 1) variety selection, 2) row spacing, 3) planting date, 4) fertility, 5) weed control, etc. Within the growing season, pests will be monitored and control provided as outlined in the objectives listed below.

Objective 1: Determine efficacy of higher use rates of Dimilin insecticide in providing season long control of pests.

Objective 2: Evaluate the effect of plant bug control on soybean yield using insecticide (Transform) application and timing strategies.

Plant bugs are generally overlooked in soybean production. They are considered a non-threat by most entomologists, but they are a major pest in cotton.

Objective 3: Evaluate the effect of plant growth regulators (Pix, Apogee, and Stance) on soybean physiology and yield.

Our goal is to identify which products/ rates/ timings will influence yields. Timings will be at various growth stages, but may be triggered based on insect populations.

Objective 4: Evaluate combinations of treatments listed in Objectives 1-3 for synergism and opportunities to minimize trips across the field.

Our desire is to identify opportunities for combining insecticides/ growth regulators to minimize trips across the field.

REPORT OF ACTIVITY/ACCOMPLISHMENTS

Objective 1: Determine efficacy of higher use rates of Dimilin insecticide in providing season long control of pests.

Throughout the two years of research Dimilin was not evaluated with growth regulator applications. This was due to Dimilin having zero effect on plant bugs and also not being a suitable tank mix partner with growth regulators. Because this objective was not included, we added an additional test evaluating application timings of Mepstar 6X(PGR) in 2020.

Objective 2: Evaluate the effect of plant bug control on soybean yield using insecticide (Transform and Orthene) application and timing strategies.

All treatments were initiated in the delta and hill locations; however, no plant bug pressure materialized at any location in 2020 or 2021.

Objective 3: Evaluate the effect of plant growth regulators (Mepstar (Pix), Stance, and Apogee) on soybean physiology and yield.

Final height and node counts were recorded between the R6 and R7 growth stages (2020 Tables 1-5) and (2021 Tables 6-8). This data revealed that plant growth regulators had no effect on total height of soybean. While not statistically significant, we did observe variation between the various growth

regulators. They did reduce plant height when compared to the untreated check, but were very inconsistent. All other locations when averaged together, height and node counts were not different from the untreated. Harvest was conducted with an Almaco SPC 40 plot combine, yields were calculated and reported in dry bu/ac with a standard moisture of 13%. Tables 1-8 show that there were no differences in yield when compared to the untreated control.

An additional experiment was performed examining the effect of multiple applications of Pix on soybean growth, development, and yield. These treatments are listed in Table 5 showing initiation at V4 growth stage and applications every two weeks following. There were single applications at every timing along with sequential treatments. Even with 4 sequential applications of Mepstar 6X, final soybean height, node, and yield exhibited no difference when compared to the untreated control (Table 5).

Objective 4: Evaluate combinations of treatments listed in Objectives 2&3 for synergism and opportunities to minimize trips across the field.

Due to low insect pressure and the lack of a growth regulator effect, no synergism/interaction was observed at any location over the two years of this study. Since no statistical differences were observed over the two years it was decided to not continue this trial for a third year.

Table 1. Effect of production inputs on soybean height, node, and yield. Chatham, Mississippi 2020

Treatment ^{ab}	Timing	Height (in) ^c	Node ^d	Yield (bu/ac) ^e
Untreated		31.5 ab	14.8 b	81.0 abc
Mepstar 6X	R1	29.9 b	15.3 ab	78.0 bc
Apogee	R1	31.2 ab	15.4 ab	88.2 abc
Stance	R1	30.1 b	15.2 ab	81.7 abc
Transform	R1	31.4 ab	15.0 b	74.9 c
Mepstar 6X	R1	31.2 ab	15.5 ab	80.3 abc
Transform	R1			
Apogee	R1	32.3 ab	15.5 ab	86.6 abc
Transform	R1			
Stance	R1	33.8 a	16.0 ab	85.8 abc
Transform	R1			
Mepstar 6X	R1	32.4 ab	15.6 ab	82.0 abc
Mepstar 6X	R3			
Apogee	R1	31.3 ab	15.6 ab	87.6 abc
Apogee	R3			
Stance	R1	31.9 ab	15.3 ab	88.7 abc
Stance	R3			
Mepstar 6X	R1	33.0 a	14.8 b	92.8 a
Transform	R1			
Mepstar 6X	R3			
Apogee	R1	32.3 ab	14.8 b	85.3 abc
Transform	R1			
Apogee	R3			
Stance	R1	31.8 ab	16.3 a	91.3 ab
Transform	R1			
Stance	R3			

Means followed by same letter or symbol do not significantly differ ($\alpha=0.05$)

^a Treatment rates at all applications: Mepstar 6X: 2 fl oz/ac; Apogee: 7.25 oz/ac; Stance: 4 fl oz/ac; Transform: 2 oz/ac

^b All Apogee applications were mixed with AMS: 1 lb/ac + COC: 1 qt/ac

^c Total plant height measured in inches (in) at growth stage R6

^d Total number of nodes counted at growth stage R6

^e Yield reported in bushels per acre (bu/ac)

Table 2. Effect of production inputs on soybean height, node, and yield. Wayside, Mississippi 2020

Treatment ^{ab}	Timing	Height (in) ^c	Node ^d	Yield (bu/ac) ^e
Untreated		37.6 abc	19.7 a	99.2 ab
Mepstar 6X	R1	39.3 a	20.1 a	103.9 ab
Apogee	R1	36.8 abc	19.7 a	94.2 b
Stance	R1	38.7 ab	20.1 a	107.8 a
Transform	R1	37.4 abc	19.5 a	93.2 b
Mepstar 6X	R1	37.4 abc	19.5 a	101.8 ab
Transform	R1			
Apogee	R1	39.2 a	19.6 a	98.9 ab
Transform	R1			
Stance	R1	36.1 bc	20.1 a	93.4 b
Transform	R1			
Mepstar 6X	R1	38.0 abc	20.3 a	100.9 ab
Mepstar 6X	R3			
Apogee	R1	39.0 a	20.4 a	102.0 ab
Apogee	R3			
Stance	R1	35.5 c	19.0 a	98.3 ab
Stance	R3			
Mepstar 6X	R1	35.8 c	19.3 a	93.6 b
Transform	R1			
Mepstar 6X	R3			
Apogee	R1	37.3 abc	19.5 a	92.6 b
Transform	R1			
Apogee	R3			
Stance	R1	35.4 c	19.7 a	100.1 ab
Transform	R1			
Stance	R3			

Means followed by same letter or symbol do not significantly differ ($\alpha=0.05$)

^a Treatment rates at all applications: Mepstar 6X: 2 fl oz/ac; Apogee: 7.25 oz/ac; Stance: 4 fl oz/ac; Transform: 2 oz/ac

^b All Apogee applications were mixed with AMS: 1 lb/ac + COC: 1 qt/ac

^c Total plant height measured in inches (in) at growth stage R6

^d Total number of nodes counted at growth stage R6

^e Yield reported in bushels per acre (bu/ac)

Table 3. Effect of production inputs on soybean height, node, and yield. Mantee, Mississippi 2020

Treatment ^{ab}	Timing	Height (in) ^c	Node ^d	Yield (bu/ac) ^e
Untreated		44.4 ab	21.4 abc	72.3 ab
Mepstar 6X	R1	41.5 b	21.3 abc	65.7 b
Apogee	R1	44.4 ab	20.9 bc	74.2 ab
Stance	R1	43.4 ab	21.7 ab	75.9 ab
Transform	R1	44.5 ab	21.4 abc	76.2 a
Mepstar 6X	R1	44.6 ab	21.8 ab	76.7 a
Transform	R1			
Apogee	R1	45.6 a	21.2 abc	74.5 ab
Transform	R1			
Stance	R1	43.7 ab	21.3 abc	72.3 ab
Transform	R1			
Mepstar 6X	R1	41.9 b	20.2 c	66.6 ab
Mepstar 6X	R3			
Apogee	R1	45.4 a	21.7 ab	75.0 ab
Apogee	R3			
Stance	R1	41.9 b	21.4 abc	72.9 ab
Stance	R3			
Mepstar 6X	R1	44.4 ab	21.0 bc	76.3 a
Transform	R1			
Mepstar 6X	R3			
Apogee	R1	44.7 ab	21.2 abc	68.5 ab
Transform	R1			
Apogee	R3			
Stance	R1	44.1 ab	22.3 a	74.8 ab
Transform	R1			
Stance	R3			

Means followed by same letter or symbol do not significantly differ ($\alpha=0.05$)

^a Treatment rates at all applications: Mepstar 6X: 2 fl oz/ac; Apogee: 7.25 oz/ac; Stance: 4 fl oz/ac; Transform: 2 oz/ac

^b All Apogee applications were mixed with AMS: 1 lb/ac + COC: 1 qt/ac

^c Total plant height measured in inches (in) at growth stage R6

^d Total number of nodes counted at growth stage R6

^e Yield reported in bushels per acre (bu/ac)

Table 4. Effect of production inputs on soybean height, node, and yield averaged across three locations in Mississippi 2020

Treatment ^{ab}	Timing	Height (in) ^c	Node ^d	Yield (bu/ac) ^e
Untreated		37.8 abc	18.7 abc	84.15 ab
Mepstar 6X	R1	36.9 bc	18.9 abc	82.54 ab
Apogee	R1	37.4 abc	18.7 abc	85.53 ab
Stance	R1	37.4 abc	19.0 abc	88.44 a
Transform	R1	37.7 abc	18.6 bc	81.98 ab
Mepstar 6X	R1	37.7 abc	19.0 abc	86.29 ab
Transform	R1			
Apogee	R1	39.0 a	18.8 abc	86.67 ab
Transform	R1			
Stance	R1	37.9 abc	19.1 abc	83.21 ab
Transform	R1			
Mepstar 6X	R1	37.4 abc	18.7 abc	80.71 b
Mepstar 6X	R3			
Apogee	R1	38.5 ab	19.2 ab	88.21 a
Apogee	R3			
Stance	R1	36.4 c	18.6 bc	86.59 ab
Stance	R3			
Mepstar 6X	R1	37.8 abc	18.3 c	87.56 ab
Transform	R1			
Mepstar 6X	R3			
Apogee	R1	38.1 abc	18.5 bc	82.11 ab
Transform	R1			
Apogee	R3			
Stance	R1	37.1 bc	19.4 a	88.73 a
Transform	R1			
Stance	R3			

Means followed by same letter or symbol do not significantly differ ($\alpha=0.05$)

^a Treatment rates at all applications: Mepstar 6X: 2 fl oz/ac; Apogee: 7.25 oz/ac; Stance: 4 fl oz/ac; Transform: 2 oz/ac

^b All Apogee applications were mixed with AMS: 1 lb/ac + COC: 1 qt/ac

^c Total plant height measured in inches (in) at growth stage R6

^d Total number of nodes counted at growth stage R6

^e Yield reported in bushels per acre (bu/ac)

Table 5. Effect of multiple applications of Mepstar 6X^a on soybean height, node, and yield.
Mantee, Mississippi 2020

Timing ^b	Height (in) ^c	Node ^d	Yield (bu/ac) ^e
Untreated	47.8 a	21.5 ab	86.8 a
V4	49.6 a	21.8 ab	87.7 a
2 Weeks	48.8 a	21.7 ab	86.6 a
4 Weeks	48.2 a	21.5 ab	86.3 a
6 Weeks	47.1 a	20.7 a	83.6 a
V4			
2 Weeks	47.9 a	21.0 ab	88.3 a
4 Weeks			
6 Weeks			
2 Weeks	48.9 a	22.3 a	87.3 a
4 Weeks			
6 Weeks			
4 Weeks	49.7 a	21.4 ab	91.7 a
6 Weeks			

Means followed by same letter or symbol do not significantly differ ($\alpha=0.05$)

^a Rate of Mepstar 6X at all application timings: 2.6 fl oz/ac

^b Timings 2 weeks, 4 weeks, 6 weeks indicate time after V4 growth stage

^c Total plant height measured in inches (in) at growth stage R6

^d Total number of nodes counted at growth stage R6

^e Yield reported in bushels per acre (bu/ac)

Table 6. Effect of production inputs on soybean height, node, and yield. Chatham, Mississippi
East 2021

Treatment ^{ab}	Timing	Height (in) ^c	Node ^d	Yield (bu/ac) ^e
Untreated		36.9a	17.9a	67.3a
Mepstar 6X	R1	34.4de	17.6a	69.9a
Apogee	R1	35.7ad	16.6a	67.8a
Stance	R1	34.0e	17.2a	67.7a
Transform	R1	35.4be	17.6a	71.9a
Mepstar 6X	R1	36.3c	18.0a	66.7a
Transform	R1			
Apogee	R1	36.6b	18.0a	62.3a
Transform	R1			
Stance	R1	34.8de	17.4a	61.5a
Transform	R1			
Mepstar 6X	R1	34.0e	17.4a	64.6a
Mepstar 6X	R3			
Apogee	R1	35.0cde	17.2a	53.8a
Apogee	R3			
Stance	R1	35.0cde	17.4a	63.3a
Stance	R3			
Mepstar 6X	R1	34.8cde	17.4a	69.3a
Transform	R1			
Mepstar 6X	R3			
Apogee	R1	35.1cde	17.2a	62.8a
Transform	R1			
Apogee	R3			
Stance	R1	34.9be	17.8a	67.5a
Transform	R1			
Stance	R3			

Means followed by same letter or symbol do not significantly differ ($\alpha=0.05$)

^a Treatment rates at all applications: Mepstar 6X: 2 fl oz/ac; Apogee: 7.25 oz/ac; Stance: 4 fl oz/ac; Transform: 2 oz/ac

^b All Apogee applications were mixed with AMS: 1 lb/ac + COC: 1 qt/ac

^c Total plant height measured in inches (in) at growth stage R6

^d Total number of nodes counted at growth stage R6

^e Yield reported in bushels per acre (bu/ac)

Table 7. Effect of production inputs on soybean height, node, and yield. Chatham, Mississippi
West 2021

Treatment ^{ab}	Timing	Height (in) ^c	Node ^d	Yield (bu/ac) ^e
Untreated		35.9a	16.1a	51.5a
Mepstar 6X	R1	35.5a	16.0a	57.7a
Apogee	R1	37.0a	15.3a	59.3a
Stance	R1	35.7a	17.3a	46.8a
Transform	R1	37.7a	15.9a	64.9a
Mepstar 6X	R1	35.6a	15.6a	64.0a
Transform	R1			
Apogee	R1	35.5a	14.8a	54.8a
Transform	R1			
Stance	R1	35.0a	15.0a	63.2a
Transform	R1			
Mepstar 6X	R1	36.2a	15.8a	64.5a
Mepstar 6X	R3			
Apogee	R1	35.9a	15.1a	54.5a
Apogee	R3			
Stance	R1	35.0a	15.5a	61.9a
Stance	R3			
Mepstar 6X	R1	35.5a	14.9a	61.8a
Transform	R1			
Mepstar 6X	R3			
Apogee	R1	35.3a	15.4a	62.2a
Transform	R1			
Apogee	R3			
Stance	R1	34.8a	15.3a	63.3a
Transform	R1			
Stance	R3			

Means followed by same letter or symbol do not significantly differ ($\alpha=0.05$)

^a Treatment rates at all applications: Mepstar 6X: 2 fl oz/ac; Apogee: 7.25 oz/ac; Stance: 4 fl oz/ac; Transform: 2 oz/ac

^b All Apogee applications were mixed with AMS: 1 lb/ac + COC: 1 qt/ac

^c Total plant height measured in inches (in) at growth stage R6

^d Total number of nodes counted at growth stage R6

^e Yield reported in bushels per acre (bu/ac)

Table 8. Effect of production inputs on soybean height, node, and yield. Mantee, Mississippi 2021

Treatment ^{ab}	Timing	Height (in) ^c	Node ^d	Yield (bu/ac) ^e
Untreated		42.5ab	18.8a	72.9ab
Mepstar 6X	R1	41.8ab	18.5a	76.9a
Apogee	R1	42.5ab	18.8a	75.4ab
Stance	R1	42.5ab	18.8a	68.1b
Transform	R1	42.3ab	18.5a	75.4ab
Mepstar 6X	R1	42.5ab	19.0a	74.9ab
Transform	R1	42.5ab	19.0a	74.9ab
Apogee	R1	42.0ab	18.8a	72.7ab
Transform	R1	42.0ab	18.8a	72.7ab
Stance	R1	41.0ab	18.0a	72.5ab
Transform	R1	41.0ab	18.0a	72.5ab
Mepstar 6X	R1	42.3ab	18.5a	73.0ab
Mepstar 6X	R3	42.3ab	18.5a	73.0ab
Apogee	R1	43.8a	19.0a	72.7ab
Apogee	R3	43.8a	19.0a	72.7ab
Stance	R1	42.3ab	18.8a	68.6b
Stance	R3	42.3ab	18.8a	68.6b
Mepstar 6X	R1	42.5ab	18.8a	72.2ab
Transform	R1	42.5ab	18.8a	72.2ab
Mepstar 6X	R3	42.5ab	18.8a	72.2ab
Apogee	R1	43.3a	19.8a	72.8ab
Transform	R1	43.3a	19.8a	72.8ab
Apogee	R3	43.3a	19.8a	72.8ab
Stance	R1	41.8ab	19.0a	70.3ab
Transform	R1	41.8ab	19.0a	70.3ab
Stance	R3	41.8ab	19.0a	70.3ab

Means followed by same letter or symbol do not significantly differ ($\alpha=0.05$)

^a Treatment rates at all applications: Mepstar 6X: 2 fl oz/ac; Apogee: 7.25 oz/ac; Stance: 4 fl oz/ac; Transform: 2 oz/ac

^b All Apogee applications were mixed with AMS: 1 lb/ac + COC: 1 qt/ac

^c Total plant height measured in inches (in) at growth stage R6

^d Total number of nodes counted at growth stage R6

^e Yield reported in bushels per acre (bu/ac)