

# Advancing Our Understanding of Charcoal Rot in Soybeans

Martha P. Romero Luna,<sup>1</sup> Daren Mueller,<sup>2</sup> Alemu Mengistu,<sup>3</sup> Asheesh K. Singh,<sup>4</sup> Glen L. Hartman,<sup>5</sup> and Kiersten A Wise<sup>1,6</sup>

<sup>1</sup>Department of Botany and Plant Pathology, Purdue University, West Lafayette, IN 47907 (mromerol@purdue.edu; kawise@purdue.edu), <sup>2</sup>Department of Plant Pathology and Microbiology, Iowa State University, Ames, IA 50011 (dsmuelle@iastate.edu), <sup>3</sup>Crop Genetics Research Unit, United States Department of Agriculture, Agricultural Research Service (USDA-ARS), 605 Airways Blvd., Jackson, TN 38301 (alemu.mengistu@ars.usda.gov), <sup>4</sup>Department of Agronomy, Iowa State University, Ames, IA 50011 (aksingh@iastate.edu), and <sup>5</sup>USDA, ARS, Department of Crop Science, University of Illinois, Urbana, IL 61801 (ghartman@illinois.edu) and <sup>6</sup>Corresponding author, e-mail: kawise@purdue.edu

The use of trade, firm, or corporation names in this paper is for the information and convenience of the reader. Such use does not constitute an official endorsement of approval by the USDA-ARS, NAL, or BIC of any product or service to the exclusion of others that may be suitable.

Subject Editor: Erin Hodgson

Received 11 October 2016; Editorial decision 19 December 2016

## Abstract

Charcoal rot [*Macrophomina phaseolina* (Tassi) Goid] of soybean [*Glycine max* (L.) Merr.] is an important but commonly misidentified disease, and very few summary articles exist on this pathosystem. Research conducted over the past 10 yr has improved our understanding of the environment conducive to disease development, host resistance, and improved disease diagnosis and management. This article summarizes the currently available research with an emphasis on disease management.

**Key words:** charcoal rot, *Macrophomina phaseolina*, soybean

## Impact of Charcoal Rot on United States Soybean Production


*Macrophomina phaseolina* (Tassi) Goid (Goidanish 1947) causes charcoal rot in soybeans [*Glycine max* (L.) Merr.], and infects ~500 other plant species including agronomic crops, fruits, legumes, and trees (Dhingra and Sinclair 1977, Dhingra and Sinclair 1978, Kunwar et al. 1986). Charcoal rot in soybean was first observed in the United States in 1949 (Young 1949). Since then, *M. phaseolina* has been observed across the United States (Wyllie and Scott 1988), causing significant yield losses in Kansas and the southern United States (Bowen and Schapaugh 1989, Wrather and Koenning 2009). Although yield losses due to charcoal rot vary among years, charcoal rot has ranked among the most important soybean diseases in the United States over the past 20 yr. In 2003 and 2012, charcoal rot was ranked as the second most important soybean disease in the United States, with an estimated yield loss of ~1.9 million and 2.0 million metric tons, respectively (Wrather et al. 2010, Allen et al. unpublished data). From 2010 to 2013, charcoal rot was among the top five yield loss-causing soybean diseases in the United States (Allen et al. unpublished data).

Although charcoal rot incidence has increased in the past 30 yr in the northern United States (Birrenkott et al. 1984, Bradley and

Río 2003, ElAraby et al. 2003, Yang and Navi 2005, Cummings and Bergstrom 2013), most of the field-based research on this disease has been conducted in the southern United States due to its wider prevalence and annual importance in this region (Gangopadhyay et al. 1970, Shokes et al. 1977, Short et al. 1978, Bowen and Schapaugh 1989, Mengistu et al. 2007).

## Disease Life Cycle and Pathogen Biology

*Macrophomina phaseolina* survives in the soil as hard, black fungal structures, called microsclerotia (Fig. 1; Crous et al. 2006). Microsclerotia germinate between 20° and 40°C, and infect root tissue. The fungus can infect seedlings and young plants as well as mature plants (Short et al. 1978, Bristow and Wyllie 1986, Collins et al. 1991). *Macrophomina phaseolina* produces enzymes and toxins that degrade root tissue, and once infection has occurred, the root and stem tissue is colonized within 2–3 wk (Islam et al. 2012). The extracellular enzymes include amylase, cellulose, hemicellulose, lipase, and pectinase (Radha 1953, Dhingra et al. 1974, Ahmad et al. 2006, Kaur 2012). Although they have been associated with disruption of the cell and membrane wall, the main activity of these extracellular enzymes, their correlation with fungal growth, and



**Fig. 1.** *Macrophomina phaseolina* microsclerotia on stem, pith, and root tissue on soybean stem.

their influence on the infection process are still largely unknown (Schinke and Germani 2012). Two toxins isolated from *M. phaseolina* that are thought to be involved in the infection process are phaseolina and – (-) botryodiplodin. Phaseolina was first identified by Siddiqui et al. (1979), and later described in more detail by Dhar et al. (1982). However, Ramezani et al. (2007) was only able to isolate – (-) botryodiplodin from *M. phaseolina* cultures, but not phaseolina. Further research is needed to confirm which toxins are produced by *M. phaseolina* and how they facilitate infection.

Under conducive environmental conditions, *M. phaseolina* will infect the vascular system, interfering with the normal plant function of transporting water and nutrients to the leaves, causing the common disease symptoms of wilting and premature leaf death (Gupta and Chauhan 2005, Khan 2007). After soybean plants are harvested, microsclerotia in the plant residue return to the soil. It has not been widely studied how long these microsclerotia survive in soil under field conditions. According to Short et al. (1980), *M. phaseolina* survives for almost 2 yr on residue at or below the soil surface; however, this study did not assess *M. phaseolina*'s ability to cause infection after this length of time. Research conducted in Brazil indicated that although microsclerotia could survive for almost 35 mo on infected crop residue, pathogenicity decreased after 6 mo (Fig. 2; Reis et al. 2014).

*Macrophomina phaseolina* is an Ascomycete, and classified in the family Botryosphaeriaceae. The sexual state of *M. phaseolina* has not yet been identified. Pycnidia production has been observed occasionally in different host plants (Dhingra and Sinclair 1978) and in artificial media such as potato dextrose agar (Gaetán et al. 2006) and water-agar-leaf medium (Chidambaram and Mathur 1975, Ma et al. 2010). The role of pycnidia on disease development is not well understood. Conidia are hyaline, ellipsoid to ovoid with dimensions of 14 to 30 × 5 to 10 μm (Crous et al. 2006). Conidia have been shown to infect soybean seedlings under laboratory conditions (Ma et al. 2010), but the role of conidia in the disease cycle in soybean is unknown.

To date there have been no reports of races or specific pathotypes within *M. phaseolina* on soybean. Many studies have been conducted trying to classify variation among isolates; however, only differences in isolate pathogenicity and virulence have been observed (Dhingra and Sinclair 1973, Mihail and Taylor 1995), and recently on regionally adapted cultivars (Sexton et al. 2016). In 1973, Dhingra and Sinclair evaluated the pathogenicity of *M. phaseolina* isolates obtained from stem, petioles, pods, and seed of three different soybean plants,


and found that isolates obtained from roots were most pathogenic. Variation among isolates was also observed based on growth rate and colony type. Mihail and Taylor (1995) evaluated the pathogenicity of 114 isolates obtained from bean, spurge (*Euphorbia lathyris*), sunflower, and sorghum under controlled conditions. Although not all isolates were able to infect all crops, all isolates were capable of infection at least one host. In a more recent study, Sexton et al. (2016) evaluated and compared pathogenicity and virulence on 42 isolates collected from soybean in the northern and southern United States. Pathogenicity and virulence were determined on soybean under controlled conditions based on the length of the stem lesion and the lesion growth rate over 9 d after inoculation, respectively. It was observed that northern isolates were more virulent compared to southern isolates on southern cultivars.

Studies on host-specificity of *M. phaseolina* have been inconclusive. Su et al. (2001) did not observe any variation after analyzing *M. phaseolina* isolates obtained from different crops with restriction fragment length polymorphism and random amplified polymorphic DNA markers. Similar findings were reported by Vandemark et al. (2000). They determined that *M. phaseolina* is a heterogeneous species and did not observe any differences in host specificity from isolates collected from different locations and hosts. However, several other studies have shown evidence for morphological differences among isolates from different hosts, including isolates collected from soybean and corn [*Zea mays* (L.)] (Pearson et al. 1987), soybean and sorghum [*Sorghum bicolor* (L.) Moench] (Cloud and Rupe 1991), and soybean and cotton [*Gossypium hirsutum* (L.)] (Jana et al. 2005). Studies by Arias et al. (2011) indicated that from the group of markers identified, the marker associated with pisatin demethylase-P450 had a unique allele for isolates from pumpkin [*Cucurbita pepo* (L.)] and snap bean [*Phaseolus vulgaris* (L.)], and another unique allele for soybean isolates.

In 2012, the *M. phaseolina* genome was sequenced and assembled (Islam et al. 2012). Large areas of synteny were observed with another important plant pathogen, *Fusarium oxysporum* Schltdl., which also has a wide host range (Islam et al. 2012). The authors suggested that since both pathogens infect a large number of plant species, these two pathogens may use similar pathways to infect hosts. Further studies on *M. phaseolina* genome may improve our understanding of its wide host range.

## Symptoms and Signs

Seedlings can be infected at emergence through the early vegetative stages, but symptoms are not typically observed until the R5 (beginning seed) to R7 (beginning maturity) growth stages (Fehr et al. 1971, Meyer et al. 1974, Mengistu 2015). Early symptoms include yellow leaves, reduced vigor, brown to red discoloration on roots and stems, and a general wilting appearance (Gupta and Chauhan 2005, Short et al. 1978). Later foliar symptoms include premature senescence, wilting, and premature plant death (Fig. 3). These symptoms are often confused with foliar symptoms of other diseases and disorders such as soybean cyst nematode (*Heterodera glycines* Ichinohe), drought stress, and early senescence. Plants that die prematurely due to charcoal rot will retain leaves attached to the petiole (Mengistu et al. 2007). The name charcoal rot comes from the gray or silver discoloration observed on the lower parts of the stem and the roots due to the formation of microsclerotia (Gupta and Chauhan 2005, Wrather et al. 2008, Gupta et al. 2012). In the field, charcoal rot symptoms are more frequently observed in stressed areas of the field such as field edges, compacted soils, and hillsides.



**Fig. 2.** Disease cycle of *Macrophomina phaseolina* (Smith et al. 2014). (A) *M. phaseolina* overwintering by microsclerotia in crop residue. (B) Infected roots after being in contact or in proximity with microsclerotia. (C) Fungus grows within the stem and root affecting and disrupting the vascular system. More microsclerotia are produced. (D) Abundant microsclerotia presence on lower stem and taproot tissue gives a charcoal-like appearance. Infected residue will become potential source of inoculum for next planted crop. Image credit: Iowa State University Integrated Pest Management Program.

Symptoms may be more pronounced in years with high temperatures and low soil moisture (Meyer et al. 1974, Mihail 1989). Plant maturity also influences charcoal rot severity (Mueller et al. 1985). Although the mechanism behind this influence of plant maturity on disease severity is not well understood, it is believed that early-maturing plants that flower during cooler temperatures will be less susceptible to high temperature stress that contributes to symptom expression and yield loss (Mihail 1989, Smith and Carvil 1997). Until recently, the association of zone lines (thin black lines in the stem cortex) was considered a symptom of charcoal rot, confusing disease diagnosis (Fig. 4). These lines are actually caused by *Diaporthe longicolla*, and not associated with charcoal rot (Olsen et al. 2015).

### Charcoal Rot Management

Although many studies have been conducted to improve disease management strategies for charcoal rot, farmers currently have limited options to minimize charcoal rot damage.

### Crop Rotation

Based on the wide host range of *M. phaseolina*, crop rotation is not considered as a reliable management practice. Short et al. (1980) found that there was a direct correlation between the population of viable microsclerotia in soil and the number of consecutive years of corn or soybean production, regardless of rotation. However, long-term crop rotation studies observed that even if soybean is rotated with another host of *M. phaseolina*, a long-term rotation will reduce the *M. phaseolina* population compared to continuous soybean planting (Francl et al. 1988, Singh et al. 1990a). It is possible that crop rotation could improve charcoal rot management if long-term rotations including nonhosts are established (Table 1); however, further research is required to confirm the impact of crop rotation on charcoal rot.

### Tillage

Baird et al. (2003) indicated that tillage may reduce charcoal rot severity by accelerating decomposition of microsclerotia or increasing antagonistic interactions. Olanya and Campbell (1988) found that





**Fig. 3.** Foliar symptoms of charcoal rot on soybean seen at the front of the image include premature senescence and plant death while leaves remain attached to the petioles.



**Fig. 4.** Zone lines (top) are a sign of diseases caused by *Diaporthe longicolla* and not associated with charcoal rot, which is characterized by the formation of microsclerotia in root and stem tissue (bottom).

one tillage pass to a depth of 15 cm reduced *M. phaseolina* populations; however, the efficacy of this practice will vary according to the initial quantity of microsclerotia present in the soil. In a later study, [Wrather et al. \(1998\)](#) observed no differences on charcoal rot

infection in soybean regardless of whether a conventional tillage or conservation tillage system was implemented, and concluded that tillage did not reduce *M. phaseolina* infection. Most recently, [Mengistu et al. \(2009\)](#) reported that colony-forming units of *M. phaseolina* in soybean tissue were greater under tillage than in no-tillage and suggested that charcoal rot may be better managed in a no-tillage system. Overall, the impact of tillage on charcoal rot has been inconsistent, and it has been difficult to establish tillage as a reliable practice for charcoal rot management.

#### Irrigation

Although *M. phaseolina* can infect under a wide range of environmental conditions, yield loss and high disease severity are often linked to areas where soils are dry or drought occurs throughout the reproductive stages of soybean growth ([Mengistu et al. 2011a](#)). In these situations, irrigating soybean to reduce drought stress can limit yield losses due to charcoal rot, even though losses may still range from 6 to 30%. In 2000, [Kendig et al.](#) observed that although disease still occurs under irrigation, it can be limited by water management. In this study, *M. phaseolina*-infected plants under water stress at the beginning of the season had higher disease severity compared with infected plants that were not water stressed early in the season. More recently, [Mengistu et al. \(2011a\)](#), observed that although infection occurred under both irrigated and nonirrigated

**Table 1.** List of several common agricultural crop hosts of *M. phaseolina*

Common name	Latin name	Reference
Corn	<i>Zea mays</i> L.	Livingston 1945
Sorghum	<i>Sorghum bicolor</i> L. Moench	Livingston 1945
Soybean	<i>Glycine max</i> L. Merr	Young 1949
Chickpea	<i>Cicer arietinum</i> L.	Dhingra and Sinclair 1977
Cotton	<i>Gossypium hirsutum</i> L.	Dhingra and Sinclair 1977
Peanut	<i>Arachis hypogaea</i> L.	Dhingra and Sinclair 1977
Sunflower	<i>Helianthus annuus</i> L.	Yang and Owen 1982, Khan 2007
Snapbean	<i>Phaseolus vulgaris</i> L.	Arias et al. 2011

environments, higher yields were reported under irrigated environments. Therefore, limiting water stress through scheduled irrigation throughout the season may reduce yield loss due to charcoal rot.

### Fertility and Plant Nutrition

There are few studies that provide information on the interaction of soil fertility on the severity of charcoal rot on soybeans. Mengistu et al. (2016) studied the effect of phosphorus and potassium on the severity of charcoal rot, and showed that *M. phaseolina* populations were not significantly impacted by phosphorus and potassium applications in no-till soils, and were not sensitive to routine phosphorus and potassium fertilizer application based on soil testing. The study suggested that soybean farmers should still apply phosphorus and potassium based on crop need without regard to potential effects of charcoal rot severity.

### Weed Control

Competition for water and other resources can stress soybean plants. Therefore, implementing recommended weed management programs is key to reducing the number of factors that contribute to stress in the soybean crop (Mengistu et al. 2015). Several studies have been conducted under greenhouse and field conditions to determine the effect of herbicides on stem and root colonization of soybeans by *M. phaseolina* (Edgar Filho and Dhingra 1980, Cerkauskas et al. 1982, Canaday et al. 1986). Some herbicides affect colonization of soybean plants by *M. phaseolina*; however, glyphosate did not affect *M. phaseolina* colonization (Canaday et al. 1986). Mengistu et al. (2013a) observed that glyphosate inhibited *M. phaseolina* growth in vitro, but under field conditions disease severity was only reduced when single glyphosate applications were performed on soybeans at early to mid-vegetative stages (V3 and V6) in a tilled environment.

### Soybean Cyst Nematode Management

The role of *H. glycines*, the cause of soybean cyst nematode, on charcoal rot development is variable. One study indicated that *H. glycines* increases root colonization by *M. phaseolina* (Todd et al. 1987), but others observed no interaction under field conditions when both organisms were present (Francl et al. 1988). Smith and Carvil (1997) did not observe an increase in either disease on soybean cyst nematode-resistant or susceptible genotypes when screened in the presence of *M. phaseolina*. In many areas of the United States, it is important to manage both diseases by selecting cultivars resistant to both soybean cyst nematode and charcoal rot.

### Fungicides

Early research on the efficacy of fungicides for charcoal rot management have focused on the use of soil fungicides that reduced microsclerotia under field conditions (Watanabe et al. 1970, Ilyas et al. 1976, Kittle and Gray 1982); however, some older chemicals induced a phytotoxic reaction in plants (Ilyas et al. 1975). Studies on fungicide efficacy and disease management have observed a reduction on *M. phaseolina* infection on roots and seedlings when soybean seed received a treatment with captan, thiram, carbendazim, thiram + carbendazim, or thiram + carboxin (Vir et al. 1972, Singh et al. 1990b, Kumar and Singh 2000, Gupta and Chauhan 2005); however, these fungicides are not commonly used in modern commercial seed treatments. Further studies are needed to test more recent fungicide active ingredients in seed treatments against charcoal rot.

### Biocontrol

Several past studies have studied the impact of antagonistic organisms on charcoal rot. Seed treated with *Rhizobacterium* strain FPT721, a plant growth-promoting rhizobacteria, reduced disease incidence under controlled conditions (Choudhary 2011). Senthikumar et al. (2009) observed inhibition and disease reduction on *M. phaseolina* when two soybean bacterial endophytes, *Bacillus* (strain: HKA-121) and *Paenibacillus* (strain: HKA-15), were tested against *M. phaseolina* under in vitro and vivo conditions, respectively. However, these biocontrol strains are not commercially available for use in soybean production.

### Cultivar Selection

With the limited options available for farmers to manage charcoal rot, most research has focused on the development of genetic resistance to charcoal rot. These studies include the development of efficient and reliable methods to characterize resistance, as well as the development of resistant cultivars. Although only moderately resistant cultivars are currently commercially available, significant progress has been made to better understand the genetic resistance behind charcoal rot. In recent years, many studies have been conducted under controlled and field conditions to determine charcoal rot-resistant soybean cultivars. Over 865 soybean genotypes have been screened for charcoal rot resistance, and of these genotypes, 23 were identified as having moderate resistance against charcoal rot (Mengistu et al. 2007, 2011b, 2013b; Pawlowski et al. 2015). Currently, charcoal rot resistance is available in a few publicly available soybean cultivars adapted primarily to southern maturity groups (Table 2).

### Screening Methods

Early screening methods focused on identifying resistance by measuring *M. phaseolina* colonization in the entire root system (Pearson et al. 1984). Later, Smith and Carvil (1997) developed a colony-forming unit assay, where microsclerotia are quantified from stem and root. This method consisted of collecting the lower stem and taproot of arbitrarily selected plants, and plating ground, dried samples on selective media. Although these screening methods provided valuable information, results from these assays were variable and there was inconsistency among studies. In 2007, Mengistu et al. studied five different assessment methods in order to establish a reliable and consistent field method for charcoal rot screening evaluations, and proposed the use of a colony-forming unit index (CFUI) as a standard method for resistance screening. This method consisted on collecting lower stem and roots of soybean plants and



**Table 2.** Publicly available soybean lines rated as moderately resistant to charcoal rot, caused by *M. phaseolina*

Genotype	Maturity Group	Reference
DG3905	III	Mengistu et al. 2011b
Manokin	IV	Mengistu et al. 2011b
DT97-4290	IV	Paris et al. 2006
DT99-16864	V	Mengistu et al. 2007 Mengistu et al. 2011b
		Gillen et al. 2016
DT99-17483	V	Mengistu et al. 2007 Mengistu et al. 2011b
DT98-7553	V	Mengistu et al. 2007 Mengistu et al. 2011b
DT99-17554	V	Mengistu et al. 2007 Mengistu et al. 2011b

plating ground, dried samples on sterilized selective media containing PDA, rifampicin, and tergitol, and incubated for 3 d at 30 °C. After 3 d, colonies are counted and converted to CFU per gram of root and stem tissue. The CFUI is later obtained by dividing the CFU for each genotype by the CFU for the genotype that has the highest CFU in the current study. According to Mengistu et al. (2014) this method allows researchers to consistently determine reactions to *M. phaseolina* across different cultivars and environments under field conditions. In 2012, Twizeyimana et al. developed a cut-stem inoculation technique where soybean cultivars could be evaluated under greenhouse conditions in a consistent and reliable manner. Disease severity is measured by the extent of necrosis that a plant presents from the inoculation point. This technique offers the advantage of quantifying the amount of inoculum in each plant tested as well as providing a precise and less time consuming measurement of disease severity, and so far it has successfully distinguished susceptible and resistant cultivars for charcoal rot in multiple studies (Mengistu et al. 2015, Sexton 2016) under greenhouse environments.

In 2016, Sexton et al. screened northern and southern *M. phaseolina* isolates on regionally adapted cultivars under greenhouse conditions. They observed regional differences between *M. phaseolina* populations and concluded that soybean cultivars should be screened under local conditions with local *M. phaseolina* isolates (Sexton et al. 2016). Ma et al. (2010) developed a *M. phaseolina* conidia production method which consisted of soaking filter paper disks in peanut butter extract supernatant. The pretreated filter papers are placed on the surface of a soybean butter extract agar sterile plate. This conidia production method and the previous inoculation technique is proposed for evaluating resistance of soybean cultivars to different isolates. This method could facilitate cultivar screening against local *M. phaseolina* isolates that was previously suggested by Sexton et al. (2015). In 2011, a quantitative DNA assay was developed to detect *M. phaseolina* from infected tissues and soil using either TaqMan or SYBER Green (Babu et al. 2011). These assays provide a reliable and accurate method for quantifying the level of pathogen in soil or plant tissue, which is likely to aid in screening assays for genetic resistance.

The soybean genome was sequenced in 2010 (Schmutz et al. 2010). Although no information on quantitative trait loci studies or genome-wide association mapping is currently available for charcoal rot in soybean, research on common bean (Hernandez-Delgado et al. 2009) has identified three novel QTL and one QTL for charcoal rot resistance, respectively. Further research will provide needed information for breeders to develop soybean cultivars with

improved levels of charcoal rot resistance by identifying resistance alleles that could be combined in moderately resistant cultivars. Improving drought tolerance through genetic resistance could also provide new possibilities for controlling charcoal rot. The combination of these studies with the traditional screening methods will likely improve resistance to charcoal rot.

## Overall Recommendations for Disease Management and Future Research Needs

Effective charcoal rot management requires an integrated approach. Farmers who have fields with a history of charcoal rot, or are in areas prone to charcoal rot development should select cultivars with the best available commercial resistance. Cultural management practices that minimize plant stress during the reproductive growth stages will also help limit damage from charcoal rot. Planting at recommended plant populations, using adequate fertility and tillage practices, controlling weeds, and irrigating where possible can help mitigate stress in soybeans, and consequently reduce the impact of charcoal rot.

Research on charcoal rot management will improve in the coming years, as there have been advances in understanding the role of resistance genes in expression of cultivar resistance, molecular interactions between the host and the pathogen, and a better understanding of the role of environment in disease development. This information will improve our ability to screen for resistant genotypes in maturity groups adapted to each region of the United States. By advancing our knowledge of this disease, we can minimize the impact of charcoal rot on soybean yield.

## Acknowledgments

This effort was funded by the United Soybean Board project number 1520-532-5608, entitled, "Improved Management of Charcoal Rot."

## References Cited

- Ahmad, Y., A. Hameed, and A. Ghaffar. 2006. Enzymatic activity of fungal pathogens in corn. *Pakistan Journal of Botany* 38: 1305–1316.
- Arias, R. S., J. D. Ray, A. Mengistu, and B. E. Scheffler. 2011. Discriminating microsatellites from *Macrophomina phaseolina* and their potential association to biological functions. *Plant Pathology* 60: 709–718.
- Babu, B. K., S. Mesapogu, A. Sharma, S. R. Somasani, and D. K. Arora. 2011. Quantitative real-time PCR assay for rapid detection of plant and human pathogenic *Macrophomina phaseolina* from field and environmental samples. *Mycologia* 103: 466–473.
- Baird, R. E., C. E. Watson, and M. Scruggs. 2003. Relative longevity of *Macrophomina phaseolina* and associated mycobiota on residual soybean roots in soil. *Plant Disease* 87: 563–566.
- Birkenkott, G. L., A. Mengistu, and C. R. Grau. 1984. First report of charcoal rot caused by *Macrophomina phaseolina* on soybeans in Wisconsin. *Plant Disease* 68: 628.
- Bowen, C. R., and W. T. Schapaugh. 1989. Relationships among charcoal rot infection, yield, and stability estimates in soybean blends. *Crop Science* 29: 42–46.
- Bradley, C. A., and L.E.D. Río. 2003. First report of charcoal rot on soybean caused by *Macrophomina phaseolina* in North Dakota. *Plant Disease* 87: 601–601.
- Bristow, P. R., and T. D. Wyllie. 1986. *Macrophomina phaseolina*, another cause of the twin-stem abnormality disease of soybean. *Plant Disease* 70: 1152–1153.
- Canaday, C. H., D. G. Helsen, and T. D. Wyllie. 1986. Effects of herbicide-induced stress on root colonization of soybeans by *Macrophomina phaseolina*. *Plant Disease* 70: 863–866.

- Cerkauskas, R. F., O. D. Dhingra, and J. B. Sinclair. 1982. Effect of herbicides on competitive saprophytic colonization by *Macrophomina phaseolina* of soybean stems. Transactions of the British Mycological Society 79: 201–205.
- Chidambaram, P., and S. B. Mathur. 1975. Production of pycnidia by *Macrophomina phaseolina*. Transactions of the British Mycological Society 64: 165IN111-168.
- Choudhary, D. K. 2011. Plant growth-promotion (PGP) activities and molecular characterization of rhizobacterial strains isolated from soybean (*Glycine max* L. Merr) plants against charcoal rot pathogen, *Macrophomina phaseolina*. Biotechnology Letters 33: 2287–2295.
- Cloud, G. L., and J. C. Rupe. 1991. Morphological instability on a chlorate medium of isolates of *Macrophomina phaseolina* from soybean and sorghum. Phytopathology 81: 892–895.
- Collins, D. J., T. D. Wyllie, and S. H. Anderson. 1991. Biological activity of *Macrophomina phaseolina* in soil. Soil Biology and Biochemistry 23: 495–496.
- Crous, P. W., B. Slippers, M. J. Wingfield, J. Rheeder, W.F.O. Marasas, A.J.L. Philips, A. Alves, T. Burgess, P. Barber, and J. Z. Groenewald. 2006. Phylogenetic lineages in the *Botryosphaeriaceae*. Studies in Mycology 55: 235–253.
- Cummings, J. A., and G. C. Bergstrom. 2013. First report of charcoal rot caused by *Macrophomina phaseolina* in soybean in New York. Plant Disease 97: 1506–1506.
- Dhar, T. K., K.A.I. Siddiqui, and E. Ali. 1982. Structure of phaseolinone, a novel phytotoxin from *Macrophomina phaseolina*. Tetrahedron Letters 23: 5459–5462.
- Dhingra, O. D., and J. B. Sinclair. 1973. Location of *Macrophomina phaseoli* on soybean plants related to culture characteristics and virulence. Phytopathology 63: 934–936.
- Dhingra, O. D., R. W. Schneider, and J. B. Sinclair. 1974. Cellulolytic and pectolytic enzymes associated with virulent and avirulent isolates of *Macrophomina phaseolina* in vitro and in soybean seedlings. Journal of Phytopathology 80: 324–329.
- Dhingra, O. D., and J. B. Sinclair. 1977. An annotated bibliography of *Macrophomina phaseolina*, 1908–1975. Imprensa Universitária, Universidade Federal de Vicosa, Minas Gerais, BR.
- Dhingra, O. D., and J. B. Sinclair. 1978. Biology and pathology of *Macrophomina phaseolina*. Imprensa Universitária, Universidade Federal de Vicosa, Minas Gerais, BR.
- Edgar Filho, S., and O. D. Dhingra. 1980. Effect of herbicides on survival of *Macrophomina phaseolina* in soil. Transactions of the British Mycological Society 74: 61–64.
- ElAraby, M. E., J. E. Kurlle, and S. R. Stetina. 2003. First report of charcoal rot (*Macrophomina phaseolina*) on soybean in Minnesota. Plant Disease 87: 202–202.
- Fehr, W. R., C. E. Caviness, D. T. Burmood, and J. S. Pennington. 1971. Stage of development descriptions for soybeans, *Glycine max* (L.) Merr. Crop Science 11: 929–931.
- Francl, L. J., T. D. Wyllie, and S. M. Rosenbrock. 1988. Influence of crop rotation on population density of *Macrophomina phaseolina* in soil infested with *Heterodera glycines*. Plant Disease 72: 760–764.
- Gangopadhyay, S., T. D. Wyllie, and V. D. Luëdders. 1970. Charcoal rot disease of soybean transmitted by seeds. Plant Disease Reporter 54: 1088–1091.
- Goidanish, G. 1947. Revisione del genere *Macrophomina* Petrak. Species tipica: *Macrophomina phaseolina* (Tass.) Goid. nov. comb. nec. *M. phaseoli* (Maubl.). Ashby Annali della Sperimentazione Agraria 1: 449–461.
- Gillen, A. M., A. Mengistu, J. R. Smith, and R. L. Paris. 2016. Registration of DT99-16864 Soybean germplasm line with moderate resistance to charcoal rot [*Macrophomina phaseolina* (Tassi) Goid.]. Journal of Plant Registrations doi:10.3198/jpr2016.01.0002crg
- Gupta, G. K., and G. S. Chauhan. 2005. Symptoms, identification and management of soybean diseases. Technical Bulletin 10. National Research Centre for Soybean, Indore, India.
- Gupta, G. K., S. K. Sharma, and R. Ramteke. 2012. Biology, epidemiology and management of the pathogenic fungus *Macrophomina phaseolina* (Tassi) Goid with special reference to charcoal rot of soybean (*Glycine max* (L.) Merr.). Journal of Phytopathology 160: 167–180.
- Hernandez-Delgado, S., M. H. Reyes-Valdés, R. Rosales-Serna, and N. Mayek-Perez. 2009. Molecular markers associated with resistance to *Macrophomina phaseolina* (Tassi) Goid in common bean. Journal of Plant Pathology 91: 163–170.
- Ilyas, M. B., M. A. Ellis, and J. B. Sinclair. 1975. Evaluation of soil fungicides for control of charcoal rot of soybeans. Plant Disease Reporter 59: 360–364.
- Ilyas, M. B., M. A. Ellis, and J. B. Sinclair. 1976. Effect of soil fungicides on *Macrophomina phaseolina* sclerotium viability in soil and in soybean stem pieces. Phytopathology 66: 355–359.
- Islam, M. S., M. S. Haque, M. M. Islam, E. M. Emdad, A. Halim, Q.M.M. Hossen, M. Z. Hossain, B. Ahmed, S. Rahim, M. S. Rahman, et al. 2012. Tools to kill: genome of one of the most destructive plant pathogenic fungi *Macrophomina phaseolina*. BMC Genomics 13: 1–16.
- Jana, T., T. R. Sharma, and N. K. Singh. 2005. SSR-based detection of genetic variability in the charcoal rot pathogen *Macrophomina phaseolina*. Mycological Research 109: 81–86.
- Livingston, J. E. 1945. Charcoal rot of corn and sorghum. Nebraska Agricultural Experimental Station Bulletin 136.
- Kaur, S. 2012. Carbohydrate degrading enzyme production by plant pathogenic mycelia and microsclerotia isolates of *Macrophomina phaseolina* through koji fermentation. Industrial Crops and Products 36: 140–148.
- Kendig, S. R., J. C. Rupe, and H. D. Scott. 2000. Effect of irrigation and soil water stress on densities of *Macrophomina phaseolina* in soil and roots of two soybean cultivars. Plant Disease 84: 895–900.
- Khan, S. N. 2007. *Macrophomina phaseolina* as causal agent for charcoal rot of sunflower. Mycopathology 5: 11–118.
- Kittle, D. R., and L. E. Gray. 1982. Response of soybeans and soybean pathogens to soil fumigation and foliar fungicide sprays. Plant Disease 66: 213–215.
- Kumar, K., and J. Singh. 2000. Location, survival, transmission and management of seed-borne *Macrophomina phaseolina*, causing charcoal rot in soybean. Annals of Plant Protection Science 8: 44–46.
- Kunwar, I. K., T. Singh, C. C. Machado, and J. B. Sinclair. 1986. Histopathology of soybean seed and seedling infection by *Macrophomina phaseolina*. Phytopathology 76: 532–535.
- Ma, J., C. B. Hill, and G. L. Hartman. 2010. Production of *Macrophomina phaseolina* conidia by multiple soybean isolates in culture. Plant Disease 94: 1088–1092.
- Mengistu, A. 2015. Charcoal rot, pp. 67–69. In G. L. Hartman, J. B. Sinclair, and J. C. Rupe, (eds.), Compendium of soybean diseases and pests. American Phytopathological Society, St. Paul, MN.
- Mengistu, A., J. D. Ray, J. R. Smith, and R. L. Paris. 2007. Charcoal rot disease assessment of soybean genotypes using a colony-forming unit index. Crop Science 47: 2453–2461.
- Mengistu, A., K. N. Reddy, R. M. Zablotowicz, and A. J. Wrather. 2009. Propagule densities of *Macrophomina phaseolina* in soybean tissue and soil as affected by tillage, cover crop, and herbicide. Plant Health Progress doi:10.1094/PHP-2009-0130-01-RS
- Mengistu, A., J. R. Smith, J. D. Ray, and N. Bellaloui. 2011a. Seasonal progress of charcoal rot and its impact on soybean productivity. Plant Disease 95: 1159–1166.
- Mengistu, A., P. A. Arelli, J. P. Bond, G. J. Shannon, and A. J. Wrather. 2011b. Evaluation of soybean genotypes for resistance to charcoal rot. Plant Health Progress doi:10.1094/PHP-2010-0926-01-RS.
- Mengistu, A., K. N. Reddy, N. Bellaloui, E. R. Walker, and H. M. Kelly. 2013a. Effect of glyphosate on *Macrophomina phaseolina* in vitro and its effect on disease severity of soybean in the field. Crop Protection 54: 23–28.
- Mengistu, A., J. Bond, R. Nelson, J. Rupe, G. Shannon, P. Arelli, and A. Wrather. 2013b. Identification of soybean accessions resistant to *Macrophomina phaseolina* by field screening and laboratory validation. Plant Health Progress doi:10.1094/PHP-2013-0318-01-RS.
- Mengistu, A., J. D. Ray, J. R. Smith, and D. L. Boykin. 2014. Maturity effects on colony-forming units of *Macrophomina phaseolina* infection as measured using near-isogenic lines of soybeans. Journal of Crop Improvement 28: 38–56.

- Mengistu, A., X. Yin, N. Bellaloui, A. M. McClure, D. D. Tyler, and K. N. Reddy. 2016. Potassium and phosphorus have no effect on severity of charcoal rot of soybean. *Canadian Journal of Plant Pathology* 38: 174–182. doi: 10.1080/07060661.2016.1168869.
- Meyer, W. A., J. B. Sinclair, and M. N. Khare. 1974. Factors affecting charcoal rot of soybean seedlings. *Phytopathology* 64: 845–849.
- Mihail, J. D. 1989. *Macrophomina phaseolina*: Spatio-temporal dynamics of inoculum and of disease in a highly susceptible crop. *Phytopathology* 79: 848–855.
- Mihail, J. D., and S. J. Taylor. 1995. Interpreting variability among isolates of *Macrophomina phaseolina* in pathogenicity, pycnidium production, and chlorate utilization. *Canadian Journal of Botany* 73: 1596–1603.
- Mueller, J. D., B. J. Short, and J. B. Sinclair. 1985. Effects of cropping history, cultivar, and sampling date on the internal fungi of soybean roots. *Plant Disease* 69: 520–523.
- Olanya, O. M., and C. L. Campbell. 1988. Effects of tillage on the spatial pattern of microsclerotia of *Macrophomina phaseolina*. *Phytopathology* 78: 217–221.
- Olsen, T. R., A. Gebreil, A. Micijevic, C. A. Bradley, K. A. Wise, D. S. Mueller, M. I. Chilvers, and F. M. Mathew. 2015. Association of *Diaporthe longicolla* with black zone lines on mature soybean (*Glycine max* L.) plants. *Plant Health Progress* doi:10.1094/PHP-RS-15-0020.
- Paris, R. L., A. Mengistu, J. M. Tyler, and J. R. Smith. 2006. Registration of soybean germplasm line DT97-4290 with moderate resistance to charcoal rot. *Crop Science* 46: 2324–2325.
- Pawlowski, M. L., C. B. Hill, and G. L. Hartman. 2015. Resistance to charcoal rot identified in ancestral soybean germplasm. *Crop Science* 55: 1230–1235.
- Pearson, C.A.S., F. W. Schwenk, and F. J. Crowe. 1984. Colonization of soybean roots by *Macrophomina phaseolina*. *Plant Disease* 68: 1086–1088.
- Pearson, C. A. S., J. F. Leslie, and F. W. Schwenk. 1987. Host preference correlated with chlorate resistance in *Macrophomina phaseolina*. *Plant Disease* 71: 828–831.
- Radha, K. 1953. The enzymatic activity of *Macrophomina phaseoli* (Maubl.), Ashby. *Plant Science* 38: 231–234.
- Ramezani, M., W. T. Shier, H. K. Abbas, J. L. Tonos, R. E. Baird, and G. L. Sciombato. 2007. Soybean charcoal rot disease fungus *Macrophomina phaseolina* in Mississippi produces the phytotoxin (-) botryodiplodin but no detectable phaseolinone. *Journal of Natural Products* 70: 128–129.
- Reis, E. M., C. Boaretto, and A.L.D. Danelli. 2014. *Macrophomina phaseolina*: density and longevity of microsclerotia in soybean root tissues and free on the soil, and competitive saprophytic ability. *Summa Phytopathologica* 40: 128–133.
- Schmutz, J., S. B. Cannon, J. Schlueter, J. Ma, T. Mitros, W. Nelson, D. L. Hyten, Q. Song, J. J. Thelen, J. Cheng, et al. 2010. Genome sequence of the palaeopolyploid soybean. *Nature* 463: 178–183.
- Schinke, C., and J. C. Germani. 2012. Screening Brazilian *Macrophomina phaseolina* isolates for alkaline lipases and other extracellular hydrolases. *International Microbiology* 15: 1–7.
- Senthilkumar, M., K. Swarnalakshmi, V. Govindasamy, Y. K. Lee, and K. Annapurna. 2009. Biocontrol potential of soybean bacterial endophytes against charcoal rot fungus, *Rhizoctonia bataticola*. *Current Microbiology* 58: 288–293.
- Sexton, Z. F., T. J. Hughes, and K. A. Wise. 2016. Analyzing isolate variability of *Macrophomina phaseolina* from a regional perspective. *Crop Protection* 81: 9–13.
- Shokes, F. M., S. D. Lyda, and W. R. Jordan. 1977. Effect of water potential on the growth and survival of *Macrophomina phaseolina*. *Phytopathology* 67: 239–241.
- Short, G. E., T. D. Wyllie, and V. D. Ammon. 1978. Quantitative enumeration of *Macrophomina phaseolina* in soybean tissues. *Phytopathology* 68: 736–741.
- Short, G. E., T. D. Wyllie, and P. R. Bristow. 1980. Survival of *Macrophomina phaseolina* in soil and residue of soybeans. *Phytopathology* 70: 13–17.
- Siddiqui, K. A. I., A. K. Gupta, A. K. Paul, and A. K. Banerjee. 1979. Purification and properties of a heat resistant exotoxin produced by *Macrophomina phaseolina* (Tassi) Goid in culture. *Experientia* 35: 1222–1223.
- Singh, S. K., Y. L. Nene, and M. V. Reddy. 1990a. Influence of cropping systems on *Macrophomina phaseolina* population in soil. *Plant Disease* 74: 812–814.
- Singh, S. N., S. K. Srivastava, P. K. Bhargava, and M. N. Khare. 1990b. Chemical control of seedling mortality in cv. JS 72-44 and bacterial pustule in cv. Punjab-1 of soybean (*Glycine max* (L.) Merrill). *Legume Research* 13: 17–20.
- Smith, D., M. Chilvers, A. Dorrance, T. Hughes, D. Mueller, T. Niblack, and K. Wise. 2014. Charcoal rot management in the north central region. University of Wisconsin Extension Bulletin A4037.
- Smith, G. S., and O. N. Carvil. 1997. Field screening of commercial and experimental soybean cultivars for their reaction to *Macrophomina phaseolina*. *Plant Disease* 81: 363–368.
- Su, G., S. O. Suh, R. W. Schneider, and J. S. Russin. 2001. Host specialization in the charcoal rot fungus, *Macrophomina phaseolina*. *Phytopathology* 91: 120–126.
- Todd, T. C., C.A.S. Pearson, and F. W. Schwenk. 1987. Effect of *Heterodera glycines* on charcoal rot severity in soybean cultivars resistant and susceptible to soybean cyst nematode. *Journal of Nematology* 19: 35.
- Twizeyimana, M., C. B. Hill, M. Pawlowski, C. Paul, and G. L. Hartman. 2012. A cut-stem inoculation technique to evaluate soybean for resistance to *Macrophomina phaseolina*. *Plant Disease* 96: 1210–1215.
- Vandemark, G., O. Martínez, V. Pecina, and M. de Jesús Alvarado. 2000. Assessment of genetic relationships among isolates of *Macrophomina phaseolina* using a simplified AFLP technique and two different methods of analysis. *Mycologia* 656–664.
- Vir, D. S., S. Gangopadhyay, and A. Gaur. 1972. Evaluation of some systemic fungicides and antibiotics against *Macrophomina phaseolina*. *Pesticides* 6: 25–26.
- Watanabe, T., R. S. Smith, and W. C. Snyder. 1970. Populations of *Macrophomina phaseoli* in soil as affected by fumigation and cropping. *Phytopathology* 60: 1717–1719.
- Wrather, J. A., S. R. Kendig, and D. D. Tyler. 1998. Tillage effects on *Macrophomina phaseolina* population density and soybean yield. *Plant Disease* 82: 247–250.
- Wrather, J. A., J. G. Shannon, T. E. Carter, J. P. Bond, J. C. Rupe, and A.M.R. Almeida. 2008. Reaction of drought-tolerant soybean genotypes to *Macrophomina phaseolina*. *Plant Health Progress* doi:10.1094/PHP-2008-0618-01-RS.
- Wrather, J. A., and S. R. Koenning. 2009. Effects of diseases on soybean yields in the United States 1996 to 2007. *Plant Health Progress* doi:10.1094/PHP-2009-0401-01-RS.
- Wrather, A., G. Shannon, R. Balardin, L. Carregal, R. Escobar, G. K. Gupta, Z. Ma, W. Morel, D. Ploper, and A. Tenuta. 2010. Effect of diseases on soybean yield in the top eight producing countries in 2006. *Plant Health Progress* doi:10.1094/PHP2010-0125-01-RS.
- Wyllie, T. D., and D. H. Scott. 1988. Soybean diseases of the north central region. APS Press, St. Paul, MN.
- Yang, S. M., and D. F. Owen. 1982. Symptomatology and detection of *Macrophomina phaseolina* in sunflower plants parasitized by *Cylindrocopturus adspersus* larvae. *Phytopathology* 72: 819–821.
- Yang, X. B., and S. S. Navi. 2005. First report of charcoal rot epidemics caused by *Macrophomina phaseolina* in soybean in Iowa. *Plant Disease* 89: 526–526.
- Young, P. A. 1949. Charcoal rot of plants in east Texas. *Bulletin Texas Agricultural Experimental Station* No. 33.